GradeMax

Pearson

Mark Scheme (Results)

January 2018

Pearson Edexcel
International Advanced Subsidiary Level
In Further Pure Mathematics F1 (WFM01)
Paper 01

edexcel

GradeMax

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2018
Publications Code WFM01_01_1801_MS
All the material in this publication is copyright
© Pearson Education Ltd 2018

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

PEARSON EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75

2. The Edexcel Mathematics mark schemes use the following types of marks:

- M marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
- A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
- B marks are unconditional accuracy marks (independent of M marks)
- Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
- ft – follow through
- the symbol $\surd$ will be used for correct ft
- cao – correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- o.e. – or equivalent (and appropriate)
- d… or dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- $*$ The answer is printed on the paper or ag- answer given
- $\lfloor$ or d… The second mark is dependent on gaining the first mark

4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.

6. If a candidate makes more than one attempt at any question:
   - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
   - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.

7. Ignore wrong working or incorrect statements following a correct answer.

General Principles for Further Pure Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

Method mark for solving 3 term quadratic:

1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q),\ \text{where} |pq| = |c|$, leading to x = ...

$(ax^2 + bx + c) = (mx + p)(nx + q),\ \text{where} |pq| = |c| \text{ and } |mn| = |a|$, leading to x = ...

2. Formula

Attempt to use the correct formula (with values for a, b and c).

3. Completing the square

Solving $x^2 + bx + c = 0$: $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0,\ \ q \neq 0$, leading to x = ...

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \to x^{n-1}$)

2. Integration

Power of at least one term increased by 1. ($x^n \to x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.


Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

January 2018 WFM01 Further Pure Mathematics F1 Mark Scheme

Question Number	Scheme	Notes	Marks
1.	Given $f(x) = 3x^2 - \frac{5}{3\sqrt{x}} - 6$, $x > 0$ and root, α, of $f(x) = 0$ lies in the interval $[1.5, 1.6]$		
(a)	$f'(x) = 6x + \frac{5}{6}x^{-\frac{3}{2}}$	At least one of either $3x^2 \to \pm Ax$ or $-\frac{5}{3\sqrt{x}} \to \pm Bx^{-\frac{3}{2}}$ where A and B are non-zero constants.	M1
		Correct differentiation which can be simplified or un-simplified	A1
	$\left\{\alpha \simeq 1.5 - \frac{f(1.5)}{f'(1.5)}\right\} \Rightarrow \alpha \simeq 1.5 - \frac{-0.6108276349...}{9.453609212...}$	**dependent on the previous M mark** Valid attempt at Newton-Raphson using their values of $f(1.5)$ and $f'(1.5)$	dM1
	$\{\alpha = 1.564613167...\} \Rightarrow \alpha = 1.565$ (3 dp)	**dependent on all 3 previous marks** 1.565 on their first iteration (Ignore any subsequent iterations)	A1 **cso**
	Correct differentiation followed by a correct answer of 1.565 scores full marks in part (a) **Correct answer with no working scores no marks in part (a)**		**(4)**
(b)	**Either** • $\frac{\alpha - 1.5}{"0.6108276349..."} = \frac{1.6 - \alpha}{"0.3623843083..."}$ • $\frac{\alpha - 1.5}{1.6 - \alpha} = \frac{"0.6108276349..."}{"0.3623843083..."}$ • $\frac{\alpha - 1.5}{"0.6108276349..."} = \frac{1.6 - 1.5}{"0.3623843083..." + "0.6108276349..."}$	A correct linear interpolation method. Do not allow this mark if a total of one or three negative lengths are used or if either fraction is the wrong way up. This mark may be implied.	M1
	Either • $\alpha = \left(\frac{(1.6)("0.6108276349...") + (1.5)("0.3623843083...")}{"0.3623843083..." + "0.6108276349..."}\right)$ • $\alpha = 1.5 + \left(\frac{"0.6108276349..."}{"0.3623843083..." + "0.6108276349..."}\right)(0.1)$ • $\alpha = 1.5 + \left(\frac{"-0.6108276349..."}{"-0.3623843083..." + "-0.6108276349..."}\right)(0.1)$	**dependent on the previous M mark** Rearranges to make $\alpha = ...$	dM1
	$\{\alpha = 1.562764092...\} \Rightarrow \alpha = 1.563$ (3 dp)	1.563 (Ignore any subsequent iterations)	A1 **cao**
			(3)
(b) **Way 2**	$\frac{x}{"0.6108276349..."} = \frac{0.1 - x}{"0.3623843083..."} \Rightarrow x = \frac{(0.1)("0.6108276349...")}{0.9732119432...} = 0.062764092...$		
	$\alpha = 1.5 + 0.062764092...$	Finds x using a correct method of similar triangles and applies "1.5 + their x"	M1 dM1
	$\{\alpha = 1.562764092...\} \Rightarrow \alpha = 1.563$ (3 dp)	1.563	A1 **cao**
(b) **Way 3**	$\frac{0.1 - x}{"0.6108276349..."} = \frac{x}{"0.3623843083..."} \Rightarrow x = \frac{(0.1)("0.3623843083...")}{0.9732119432...} = 0.037235908...$		
	$\alpha = 1.6 - 0.037235908...$	Finds x using a correct method of similar triangles and applies "1.6 – their x"	M1 dM1
	$\{\alpha = 1.562764092...\} \Rightarrow \alpha = 1.563$ (3 dp)	1.563	A1 **cao**
			7

	Question 1 Notes	
1. (a)	**Note**	Incorrect differentiation followed by their estimate of α with no evidence of applying the NR formula is final dM0A0.
	dM1	This mark can be implied by applying at least one correct ***value*** of either $f(1.5)$ or $f'(1.5)$ to 1 significant figure in $1.5 - \frac{f(1.5)}{f'(1.5)}$. So just $1.5 - \frac{f(1.5)}{f'(1.5)}$ with an incorrect answer and no other evidence scores final dM0A0.
	Note	You can imply the M1A1 marks for algebraic differentiation for either • $f'(1.5) = 6(1.5) + \frac{5}{6}(1.5)^{-\frac{3}{2}}$ • $f'(1.5)$ applied correctly in $\alpha \simeq 1.5 - \frac{3(1.5)^2 - \frac{5}{3}(1.5)^{-\frac{1}{2}} - 6}{6(1.5) + \frac{5}{6}(1.5)^{-\frac{3}{2}}}$
	Note	**Differentiating INCORRECTLY to give** $f'(x) = 6x - \frac{5}{6}x^{-2}$ leads to $\alpha \simeq 1.5 - \frac{-0.6108276349...}{9.3703703704...} = 1.565187139... = 1.565$ (3 dp) **This response should be awarded M1 A0 dM1 A0**
	Note	**Differentiating INCORRECTLY to give** $f'(x) = 6x - \frac{5}{6}x^{-\frac{3}{2}}$ leads to $\alpha \simeq 1.5 - \frac{-0.6108276349...}{8.546390788...} = 1.571471999... = 1.571$ (3 dp) **This response should be awarded M1 A0 dM1 A0**
	S.C.	**Special Case: Differentiating INCORRECTLY to give** $f'(x) = 6x - \frac{5}{6}x^{-\frac{3}{2}}$ and $\alpha \simeq 1.5 - \frac{f(1.5)}{f'(1.5)} = 1.571$ is **M1 A0 dM1 A0**
1. (b)	**Note**	$\frac{\alpha - 1.5}{1.6 - \alpha} = \left\lvert\frac{"-0.6108276349..."}{"0.3623843083..."}\right\rvert$ is a valid method for the first M mark
	Note	$\frac{\alpha - 1.5}{1.6 - \alpha} = \frac{"0.6108276349..."}{"0.3623843083..."} \Rightarrow \alpha = 1.563$ with no intermediate working is M1 dM1 A1
	Note	$\frac{\alpha - 1.5}{-0.6108276349...} = \frac{1.6 - \alpha}{0.3623843083...} \Rightarrow \alpha = 1.745861961... = 1.745$ (3 dp) is M0 dM0 A0
	Note	$\frac{\alpha - 1.5}{-0.6108276349...} = \frac{1.6 - \alpha}{-0.3623843083...} \Rightarrow \alpha = 1.562764092... = 1.563$ (3 dp) is M1 dM1 A1

Question Number	Scheme	Notes	Marks
2.	$f(z) = z^4 - 6z^3 + 38z^2 - 94z + 221$, $z_1 = 2 + 3i$ satisfies $f(z) = 0$		
(a)	$\{z_2 =\}\ 2 - 3i$	$2 - 3i$ seen or used in part (a)	B1
	$z^2 - 4z + 13$	Attempt to expand $(z-(2+3i))(z-(2-3i))$ or $(z-(2+3i))(z-(\text{their complex } z_2))$ or any valid method ***to establish a quadratic factor*** e.g. $z = 2 \pm 3i \Rightarrow z - 2 = \pm 3i \Rightarrow z^2 - 4z + 4 = -9$ or sum of roots $= 4$, product of roots 13 to give $z^2 \pm (\text{their sum})z + (\text{their product})$	M1
		$z^2 - 4z + 13$	A1
	$(z^2 - 4z + 13)(z^2 - 2z + 17)$	Attempts to find the other quadratic factor. e.g. using long division to obtain either $z^2 \pm kz + ..., k = \text{value} \neq 0$ or $z^2 \pm \alpha z + \beta, \beta = \text{value} \neq 0, \alpha$ can be 0 or e.g. factorising to obtain either $f(z) = (z^2 - 2z + 5)(z^2 \pm kz \pm c), k = \text{value} \neq 0$ or $f(z) = (z^2 - 2z + 5)(z^2 \pm \alpha z \pm \beta), \beta = \text{value} \neq 0, \alpha$ can be 0	M1
		$z^2 - 2z + 17$	A1
	$\{z^2 - 2z + 17 = 0 \Rightarrow\}$		
	Either • $z = \dfrac{-(-2) \pm \sqrt{(-2)^2 - 4(1)(17)}}{2(1)}$ • $(z-1)^2 - 1 + 17 = 0 \Rightarrow z = ...$	**dependent on only the previous M mark** Correct method of applying the quadratic formula or completing the square for solving a 3TQ on their 2nd quadratic factor	dM1
	$\{z =\}\ 1 + 4i, 1 - 4i$	$1 + 4i$ **and** $1 - 4i$	A1
			(7)
(b)		**Criteria** • $2 \pm 3i$ plotted correctly in quadrants 1 and 4 • **Dependent on the final M mark being awarded in part (a).** Their final two roots are plotted correctly	
		Satisfies at least one of the criteria	B1ft
		Satisfies both criteria with some indication of scale or coordinates stated with at least one pair of roots symmetrical about the real axis	B1ft
			(2)
			9

	Question 2 Notes	
2. (a)	**Note**	No working leading to $x = 1+4\text{i},\ 1-4\text{i}$ is M0A0M0A0M0A0.
	Note	You can assume $x \equiv z$ for solutions in this question.
	Note	Give dM1A1 for $z^2 - 2z + 17 = 0 \Rightarrow z = 1 + 4\text{i}, 1 - 4\text{i}$ with no intermediate working.
	Note	**<u>Special Case:</u>** If their second ***3 term quadratic*** factor **can** be factorised then give Special Case dM1 for correct factorisation leading to $z = ...$
	Note	Otherwise, give 3rd dM0 for applying a method of factorising to solve their 3TQ.
	Note	**Reminder:** Method Mark for solving a 3TQ, "$az^2 + bz + c = 0$" **Formula:** Attempt to use the correct formula (with values for a, b and c) **Completing the square:** $\left(z \pm \frac{b}{2}\right)^2 \pm q \pm c = 0,\ q \neq 0,$ leading to $z = ...$

Question Number	Scheme	Notes	Marks
3. (a)	$\sum_{r=1}^{n} r^2(r+1) = \sum_{r=1}^{n} r^3 + \sum_{r=1}^{n} r^2$		
	$= \frac{1}{4}n^2(n+1)^2 + \frac{1}{6}n(n+1)(2n+1)$	Attempts to expand $r^2(r+1)$ and attempts to substitute at least one correct standard formula into their resulting expression.	M1
		Correct expression (or equivalent)	A1
	$= \frac{1}{12}n(n+1)\left[3n(n+1)+2(2n+1)\right]$	**dependent on the previous M mark** Attempt to factorise at least $n(n+1)$ having attempted to substitute both standard formulae.	dM1
	$= \frac{1}{12}n(n+1)\left[3n^2+7n+2\right]$	{this step does not have to be written}	
	$= \frac{1}{12}n(n+1)(n+2)(3n+1)$	Correct completion with no errors. **Note:** $a=3$, $b=1$	A1
			(4)
(b)	$\sum_{r=5}^{25} r^2(r+1) + \sum_{r=1}^{k} 3^r = 140543$	{**Note:** Let $f(n) = \frac{1}{12}n(n+1)(n+2)(3n+1)$ ***or their*** answer to part (a).}	
	$\left\{\sum_{r=5}^{25} r^2(r+1)\right\} = \left(\frac{1}{12}(25)(26)(27)(76)\right) - \left(\frac{1}{12}(4)(5)(6)(13)\right)$ $\{= 111150 - 130 = 111020\}$	Attempts to find either $f(25) - f(4)$ or $f(25) - f(5)$ This mark can be implied	M1
	$\sum_{r=1}^{k} 3^r = 140543 - "111020"$ $\{= 29523\}$	**dependent on the previous M mark** their $\sum_{r=1}^{k} 3^r = 140543 - "111020"$ This mark can be implied	dM1
	$\frac{3(1-3^k)}{1-3}$ or $\frac{3(3^k-1)}{3-1}$	Correct GP sum formula with $a=3$, $r=3$, $n=k$	M1
	$\left\{\frac{3(1-3^k)}{1-3} = 29523 \Rightarrow 3^k = 19683 \Rightarrow\right\}$ $k=9$	$k=9$ from a correct solution	A1 **cso**
			(4)
(b) **Alt 1**	**Alt 1 Method for the final 2 marks**		
	$\sum_{r=1}^{k} 3^r = 29523$ $\Rightarrow 3+3^2+3^3+3^4+3^5+3^6+3^7+3^8+3^9$ or $3+9+27+81+243+729+2187+6561+19683$	Attempts to solve $\sum_{r=1}^{k} 3^r =$ value by evaluating 3^r from $r=1$ to at least as far as $r=9$	M1
	$= 29523$, so $k=9$	$k=9$ from a correct solution	A1 **cso**
(b) **Alt 2**	**Alt 2 Method for the final 2 marks**		
	$\sum_{r=1}^{k} 3^r = 29523 \Rightarrow 3(1+3+3^2+3^3+...+3^{k-1}) = 29523$		
	$\left\{\sum_{r=1}^{k} 3^r = \sum_{r=1}^{k-1} 3^r + 3^k =\right\}$ $\frac{"29523"}{3} - 1 + 3^k = "29523"$	$\frac{"29523"}{3} - 1 + 3^k = "29523"$	M1
	$\{3^k = 19683 \Rightarrow\}$ $k=9$	$k=9$ from a correct solution	A1 **cso**
			8

		Question 3 Notes
3. (a)	**Note**	Applying e.g. $n=1,\ n=2$ to the printed equation without applying the standard formulae to give $a=3,\ b=1$ is M0A0M0A0
	Alt 1	**Alt Method 1 (Award the first two marks using the main scheme)**
		Using $\frac{1}{12}\left(3n^4+10n^3+9n^2+2n\right)\equiv\frac{1}{12}\left(an^4+(3a+b)n^3+(2a+3b)n^2+2bn\right)$ o.e.
	dM1	Equating coefficients to find both $a=...$ and $b=...$ **and** at least one of $a=3,\ b=1$
	A1 cso	Finds $a=3,\ b=1$ and demonstrates the identity works for all of its terms.
	Alt 2	**Alt Method 2: (Award the first two marks using the main scheme)**
		$\frac{1}{4}n^2(n+1)^2+\frac{1}{6}n(n+1)(2n+1)\equiv\frac{1}{12}n(n+1)(n+2)(an+b)$
	dM1	Substitutes $n=1,\ n=2,$ into this identity o.e. and solves to find both $a=...$ and $b=...$ **and** at least one of $a=3,\ b=1$. **Note:** $n=1$ gives $4=a+b$ and $n=2$ gives $7=2a+b$
	A1	Finds $a=3,\ b=1$
	Note	Allow final dM1A1 for $\frac{1}{4}n^4+\frac{5}{6}n^3+\frac{3}{4}n^2+\frac{1}{6}n$ or $\frac{1}{12}n(3n^3+10n^2+9n+2)$
		or $\frac{1}{12}(3n^4+10n^3+9n^2+2n)\ \rightarrow\frac{1}{12}n(n+1)(n+2)(3n+1)$ with no incorrect working.
	Note	A correct proof $\sum_{r=1}^{n}r^2(r+1)=\frac{1}{12}n(n+1)(n+2)(3n+1)$ followed by stating an incorrect
		e.g. $a=1,\ b=3$ is M1A1dM1A1 (ignore subsequent working)
(b)	**Note**	Using $\mathrm{f}(25)-\mathrm{f}(5)$ gives • $\mathrm{f}(25)-\mathrm{f}(5)=111150-280=110870$ • $\sum_{r=1}^{k}3^r=140543-"110870"=29673$
	Note	Allow 1st M1 for either • $\left\{\sum_{r=5}^{25}r^2(r+1)\right\}=\left(\frac{1}{4}(25)^2(26)^2+\frac{1}{6}(25)(26)(51)\right)-\left(\frac{1}{4}(4)^2(5)^2+\frac{1}{6}(4)(5)(9)\right)$ $\{=(105625+5525)-(100+30)=111150-130=111020\}$ • $\left\{\sum_{r=5}^{25}r^2(r+1)\right\}=\left(\frac{1}{4}(25)^2(26)^2+\frac{1}{6}(25)(26)(51)\right)-\left(\frac{1}{4}(5)^2(6)^2+\frac{1}{6}(5)(6)(11)\right)$ $\{=(105625+5525)-(225+55)=111150-280=110870\}$
	Note	$\frac{3\left(1-3^k\right)}{1-3}$ or $\frac{3\left(3^k-1\right)}{3-1}=29523\Rightarrow k=9$ with no intermediate working is 2nd M1 2nd A1
	Note	$\sum_{r=1}^{k}3^r=29523\Rightarrow k=9$ with no intermediate working is 2nd M1 2nd A1

Question Number	Scheme	Notes	Marks
4.	$3x^2+2x+5=0$ has roots α, β		
(a)	$\alpha+\beta=-\frac{2}{3},\ \alpha\beta=\frac{5}{3}$		
	$\alpha^2+\beta^2=(\alpha+\beta)^2-2\alpha\beta=.....$	**Use** of the **correct** identity for $\alpha^2+\beta^2$ (May be implied by their work)	M1
	$=\left(-\frac{2}{3}\right)^2-2\left(\frac{5}{3}\right)=-\frac{26}{9}$	$-\frac{26}{9}$ or $-2\frac{8}{9}$ from correct working	A1 **cso**
			(2)
(b)	$\alpha^3+\beta^3=(\alpha+\beta)^3-3\alpha\beta(\alpha+\beta)=.....$ or $=(\alpha+\beta)(\alpha^2+\beta^2-\alpha\beta)=.....$	**Use** of an **appropriate** and **correct** identity for $\alpha^3+\beta^3$ (May be implied by their work)	M1
	$=\left(-\frac{2}{3}\right)^3-3\left(\frac{5}{3}\right)\left(-\frac{2}{3}\right)=\frac{82}{27}$ * or $=\left(-\frac{2}{3}\right)\left(-\frac{26}{9}-\frac{5}{3}\right)=\frac{82}{27}$ *	$\frac{82}{27}$ **from correct working**	A1 * **cso**
			(2)
(c)	Sum $=\alpha+\frac{\alpha}{\beta^2}+\beta+\frac{\beta}{\alpha^2}$ $=\alpha+\beta+\frac{\alpha^3+\beta^3}{(\alpha\beta)^2}$ or $=\frac{\alpha\beta^2+\alpha}{\beta^2}+\frac{\alpha^2\beta+\beta}{\alpha^2}$ $=\frac{\alpha^3+\beta^3+\alpha^2\beta^2(\alpha+\beta)}{\alpha^2\beta^2}$ $=\left(-\frac{2}{3}\right)+\frac{\left(\frac{82}{27}\right)}{\left(\frac{5}{3}\right)^2}\ \left\{=-\frac{2}{3}+\frac{82}{75}=\frac{32}{75}\right\}$	Simplifies $\frac{\alpha}{\beta^2}+\frac{\beta}{\alpha^2}$ to give either $\frac{\alpha^3+\beta^3}{(\alpha\beta)^2}$ or $\frac{\alpha^3+\beta^3}{\alpha^2\beta^2}$ **and** substitutes at least one of their $\alpha+\beta$, $\alpha^3+\beta^3$ or $\alpha\beta$ into an expression for the sum of $\left(\alpha+\frac{\alpha}{\beta^2}\right)$ and $\left(\beta+\frac{\beta}{\alpha^2}\right)$	M1
	Product $=\left(\alpha+\frac{\alpha}{\beta^2}\right)\left(\beta+\frac{\beta}{\alpha^2}\right)$ $=\alpha\beta+\frac{\alpha\beta}{\alpha^2}+\frac{\alpha\beta}{\beta^2}+\frac{\alpha\beta}{\alpha^2\beta^2}$ $=\alpha\beta+\frac{\beta}{\alpha}+\frac{\alpha}{\beta}+\frac{1}{\alpha\beta}$ $=\alpha\beta+\frac{\beta^2+\alpha^2}{\alpha\beta}+\frac{1}{\alpha\beta}$ or $=\left(\frac{\alpha\beta^2+\alpha}{\beta^2}\right)\left(\frac{\alpha^2\beta+\beta}{\alpha^2}\right)$ $=\frac{\alpha^3\beta^3+\alpha\beta^3+\alpha^3\beta+\alpha\beta}{\alpha^2\beta^2}$ $=\frac{\alpha^3\beta^3+\alpha\beta(\beta^2+\alpha^2)+\alpha\beta}{\alpha^2\beta^2}$ $=\left(\frac{5}{3}\right)+\frac{\left(-\frac{26}{9}\right)}{\left(\frac{5}{3}\right)}+\frac{1}{\left(\frac{5}{3}\right)}\ \left\{=\frac{5}{3}-\frac{26}{15}+\frac{3}{5}=\frac{8}{15}\right\}$	Expands $\left(\alpha+\frac{\alpha}{\beta^2}\right)\left(\beta+\frac{\beta}{\alpha^2}\right)$ to give 4 terms and substitutes either their $\alpha\beta$ at least once or their $\alpha^2+\beta^2$ into their resulting expression	M1
	$x^2-\frac{32}{75}x+\frac{8}{15}=0$	Applies $x^2-(\text{sum})x+\text{product}$ (can be implied), where sum and product are numerical values. **Note:** "$=0$" not required for this mark	M1
	$75x^2-32x+40=0$	Any integer multiple of $75x^2-32x+40=0$, including the "$=0$"	A1
			(4)
			8

	Question 4 Notes	
4. (a)	**Note**	Writing a correct $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$ without attempting to substitute at least one of either their $\alpha + \beta$ or their $\alpha\beta$ into $(\alpha + \beta)^2 - 2\alpha\beta$ is M0
	Note	Give M1A0 for $\alpha + \beta = \frac{2}{3},\ \alpha\beta = \frac{5}{3}$ leading to $\alpha^2 + \beta^2 = \left(\frac{2}{3}\right)^2 - 2\left(\frac{5}{3}\right) = -\frac{26}{9}$
	Note	Give M1A1 for writing $\alpha^2 + \beta^2 = -\frac{26}{9}$ with no evidence of applying $\alpha + \beta = -\frac{2}{3},\ \alpha\beta = \frac{5}{3}$
(b)	**Note**	Allow M1 A1 for $\alpha^3 + \beta^3 = (\alpha^2 + \beta^2)(\alpha + \beta) - \alpha\beta(\alpha + \beta)$ $= \left(-\frac{26}{9}\right)\left(-\frac{2}{3}\right) - \left(-\frac{2}{3}\right)\left(\frac{5}{3}\right) \left\{= \frac{52}{27} + \frac{10}{9}\right\} = \frac{82}{27}$ *
	Note	Writing a correct $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$ without attempting to substitute at least one of either their $\alpha + \beta$ or their $\alpha\beta$ into $(\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$ is M0
	Note	Writing a correct $\alpha^3 + \beta^3 = (\alpha + \beta)(\alpha^2 + \beta^2 - \alpha\beta)$ without attempting to substitute at least one of either their $\alpha + \beta$, their $\alpha^2 + \beta^2$ or their $\alpha\beta$ into $(\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$ is M0
(a), (b)	**Note**	Applying $\frac{-1+\sqrt{14}\,i}{3}, \frac{-1-\sqrt{14}\,i}{3}$ explicitly will score (a) M0A0, (b) M0A0 • E.g. In part (a), give no credit for $\left(\frac{-1+\sqrt{14}\,i}{3}\right)^2 + \left(\frac{-1-\sqrt{14}\,i}{3}\right)^2 = -\frac{26}{9}$ • E.g. In part (b), give no credit for $\left(\frac{-1+\sqrt{14}\,i}{3}\right)^3 + \left(\frac{-1-\sqrt{14}\,i}{3}\right)^3 = \frac{82}{17}$
	Note	Using $\frac{-1+\sqrt{14}\,i}{3}, \frac{-1-\sqrt{14}\,i}{3}$ to find $\alpha + \beta = -\frac{2}{3},\ \alpha\beta = \frac{5}{3}$ followed by • $\alpha^2 + \beta^2 = \left(\frac{2}{3}\right)^2 - 2\left(\frac{5}{3}\right) = -\frac{26}{9}$, scores M1A0 in part (a) • $\alpha^3 + \beta^3 = \left(-\frac{2}{3}\right)^3 - 3\left(\frac{5}{3}\right)\left(-\frac{2}{3}\right) = \frac{82}{27}$, scores M1A0 in part (b)
(c)	**Note**	A correct method leading to $a = 75, b = -32, c = 40$ without writing a final answer of $75x^2 - 32x + 40 = 0$ is final M1A0.
	Note	Using $\frac{-1+\sqrt{14}\,i}{3}, \frac{-1-\sqrt{14}\,i}{3}$ ***explicitly*** to find the sum and product of $\left(\alpha + \frac{\alpha}{\beta^2}\right)$ and $\left(\beta + \frac{\beta}{\alpha^2}\right)$ scores M0M0M0A0 in part (c).
	Note	Using $\frac{-1+\sqrt{14}\,i}{3}, \frac{-1-\sqrt{14}\,i}{3}$ to find $\alpha + \beta = -\frac{2}{3},\ \alpha\beta = \frac{5}{3}$ **and applying** $\alpha + \beta = -\frac{2}{3},\ \alpha\beta = \frac{5}{3}$ can potentially score full marks in part (c). E.g. • Sum $= \alpha + \beta + \frac{\alpha^3 + \beta^3}{(\alpha\beta)^2} = \left(-\frac{2}{3}\right) + \frac{\left(\frac{82}{27}\right)}{\left(\frac{5}{3}\right)^2} = \frac{32}{75}$ • Product $= \alpha\beta + \frac{\beta^2 + \alpha^2}{\alpha\beta} + \frac{1}{\alpha\beta} = \left(\frac{5}{3}\right) + \frac{\left(-\frac{26}{9}\right)}{\left(\frac{5}{3}\right)} + \frac{1}{\left(\frac{5}{3}\right)} = \frac{8}{15}$ • $x^2 - \frac{32}{75}x + \frac{8}{15} = 0 \Rightarrow 75x^2 - 32x + 40 = 0$

Question Number	Scheme	Notes	Marks
5.	(i) $\dfrac{2z+3}{z+5-2\mathrm{i}} = 1+\mathrm{i}$ (ii) $w=(3+\lambda\mathrm{i})(2+\mathrm{i})$ and $\lvert w\rvert = 15$		
(i)	$2z+3=(1+\mathrm{i})(z+5-2\mathrm{i})$	Multiplies both sides by $(z+5-2\mathrm{i})$	M1
	$2z+3=z+5-2\mathrm{i}+\mathrm{i}z+5\mathrm{i}+2=z+\mathrm{i}z+7+3\mathrm{i}$		
	E.g. • $2z-z(1+\mathrm{i})=(1+\mathrm{i})(5-2\mathrm{i})-3$ • $z-\mathrm{i}z=4+3\mathrm{i}$	**dependent on the previous M mark** Collects terms in z to one side	dM1
	$z=\dfrac{4+3\mathrm{i}}{1-\mathrm{i}}$	Correct expression for $z=...$	A1
	$z=\dfrac{(4+3\mathrm{i})}{(1-\mathrm{i})}\dfrac{(1+\mathrm{i})}{(1+\mathrm{i})}=\dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$	**dependent on both previous M marks** Multiplies numerator and denominator by the conjugate of the denominator and attempts to find $z=...$	ddM1
		e.g. $\dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$ **or** $\dfrac{7}{2}\mathrm{i}+\dfrac{1}{2}$ **or** $0.5+3.5\mathrm{i}$ **or** $a=\dfrac{1}{2}, b=\dfrac{7}{2}$	A1 **cao**
			(5)
(i) **Way 2**	$2z+3=(1+\mathrm{i})(z+5-2\mathrm{i})$	Multiplies both sides by $(z+5-2\mathrm{i})$	M1
	$2(a+b\mathrm{i})+3=(1+\mathrm{i})(a+b\mathrm{i}+5-2\mathrm{i})$ $(2a+3)+2b\mathrm{i}=a+b\mathrm{i}+5-2\mathrm{i}+a\mathrm{i}-b+5\mathrm{i}+2$ $(2a+3)+2b\mathrm{i}=(a-b+7)+(b+a+3)\mathrm{i}$ $\{\text{Real}\Rightarrow\}\quad 2a+3=a-b+7$ $\{\text{Imaginary}\Rightarrow\}\quad 2b=b+a+3$	**dependent on the previous M mark** Applies $z=a+b\mathrm{i}$, multiplies out and attempts to equate **either** the real part **or** the imaginary part of the resulting equation	dM1
		Both correct equations which can be simplified or un-simplified	A1
	$\left\{\begin{array}{l} a+b=4 \\ -a+b=3 \end{array}\right\} \Rightarrow b=\dfrac{7}{2}, a=\dfrac{1}{2}$	**dependent on both previous M marks.** Obtains two equations both in terms of a and b and solves them simultaneously to give at least one of $a=...$ or $b=...$	ddM1
		e.g. $a=\dfrac{1}{2}, b=\dfrac{7}{2}$ **or** $\dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$ **or** $\dfrac{7}{2}\mathrm{i}+\dfrac{1}{2}$ **or** $0.5+3.5\mathrm{i}$	A1 **cao**
			(5)
(ii)	$w=6+3\mathrm{i}+2\mathrm{i}\lambda-\lambda$ $w=(6-\lambda)+(3+2\lambda)\mathrm{i}$ $(15)^2=(6-\lambda)^2+(3+2\lambda)^2$	Squares and adds the real and imaginary parts of w and sets equal to either 15^2 or 15	M1
		Correct equation which can be simplified or un-simplified	A1
	$\{225=36-12\lambda+\lambda^2+9+12\lambda+4\lambda^2\}$ $225=45+5\lambda^2 \Rightarrow \lambda^2=36$	**dependent on the previous M mark** Solves their quadratic in λ to give $\lambda^2=...$ or $\lambda=...$	dM1
	$\lambda=6,-6$	$\lambda=6,-6$	A1
			(4)
(ii) **Way 2**	$\{\lvert(3+\lambda\mathrm{i})(2+\mathrm{i})\rvert=15\Rightarrow\}$ $\sqrt{(3^2+\lambda^2)}\sqrt{(2^2+1^2)}=15$ or $(3^2+\lambda^2)(5)=(15)^2$	$\sqrt{(3^2+\lambda^2)}\sqrt{(2^2+1^2)}=15$ or $(3^2+\lambda^2)(2^2+1^2)=15$	M1
		Correct equation which can be simplified or un-simplified	A1
	$45=9+\lambda^2 \Rightarrow \lambda^2=36$	**dependent on the previous M mark** Solves their quadratic in λ to give $\lambda^2=...$ or $\lambda=...$	dM1
	$\lambda=6,-6$	$\lambda=6,-6$	A1
			(4)
			9

Question Number	Scheme	Notes	Marks
5.	$\dfrac{2z+3}{z+5-2\mathrm{i}} = 1+\mathrm{i}$		
(i) **Way 3**	$\dfrac{2z+10-4\mathrm{i}-7+4\mathrm{i}}{z+5-2\mathrm{i}} = 1+\mathrm{i}$		
	$2+\dfrac{-7+4\mathrm{i}}{z+5-2\mathrm{i}} = 1+\mathrm{i}$	$\dfrac{2z+3}{z+5-2\mathrm{i}} \to 2 \pm \dfrac{k}{z+5-2\mathrm{i}}$, $k \in \mathbb{C}$	M1
	$1-\mathrm{i} = \dfrac{7-4\mathrm{i}}{z+5-2\mathrm{i}}$		
	$z+5-2\mathrm{i} = \dfrac{7-4\mathrm{i}}{1-\mathrm{i}}$	**dependent on the previous M mark** Rearranges to give $z+5-2\mathrm{i} = ...$	dM1
		Correct expression for $z+5-2\mathrm{i} = ...$	A1
	$z+5-2\mathrm{i} = \dfrac{(7-4\mathrm{i})}{(1-\mathrm{i})}\dfrac{(1+\mathrm{i})}{(1+\mathrm{i})} \Rightarrow z = ...$	**dependent on both previous M marks** Multiplies numerator and denominator by the conjugate of the denominator and attempts to find $z = ...$	ddM1
	$\left\{z+5-2\mathrm{i} = \dfrac{11}{2}+\dfrac{3}{2}\mathrm{i} \Rightarrow\right\}\ z = \dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$	e.g. $\dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$ **or** $\dfrac{7}{2}\mathrm{i}+\dfrac{1}{2}$ **or** $0.5+3.5\mathrm{i}$	A1
			(5)
(i) **Way 4**	$\dfrac{2(a+b\mathrm{i})+3}{a+b\mathrm{i}+5-2\mathrm{i}} = 1+\mathrm{i} \Rightarrow \dfrac{(2a+3)+2b\mathrm{i}}{(a+5)+(b-2)\mathrm{i}} = 1+\mathrm{i}$		
	$\left(\dfrac{(2a+3)+2b\mathrm{i}}{(a+5)+(b-2)\mathrm{i}}\right)\left(\dfrac{(a+5)-(b-2)\mathrm{i}}{(a+5)-(b-2)\mathrm{i}}\right) = 1+\mathrm{i}$		
	$\dfrac{\left[(2a+3)(a+5)+2b(b-2)\right]+\mathrm{i}\left[2b(a+5)-(2a+3)(b-2)\right]}{(a+5)^2+(b-2)^2} = 1+\mathrm{i}$		
	$\{\text{Real} \Rightarrow\}\ \dfrac{(2a+3)(a+5)+2b(b-2)}{(a+5)^2+(b-2)^2} = 1$ $\{\text{Imaginary} \Rightarrow\}\ \dfrac{2b(a+5)-(2a+3)(b-2)}{(a+5)^2+(b-2)^2} = 1$	Applies $z = a+b\mathrm{i}$ and a full method leading to equating both the real part and the imaginary part	M1
	$\{\text{Real} \Rightarrow\}\ a^2+b^2+3a-14 = 0$ $\{\text{Imaginary} \Rightarrow\}\ a^2+b^2+6a-11b+23 = 0$	**dependent on the previous M mark** Manipulates both their real part and their imaginary part into their simplest forms	dM1
		Both correct simplified equations	A1
	"Real - Imaginary" gives $-3a+11b-37 = 0$ **and** e.g. • $a = \dfrac{11b-37}{3} \Rightarrow \left(\dfrac{11b-37}{3}\right)^2+b^2+3\left(\dfrac{11b-37}{3}\right)-14 = 0$ $\Rightarrow 2b^2-11b+14 = 0 \Rightarrow (b-2)(2b-7) = 0 \Rightarrow b = ...$ • $b = \dfrac{3a+37}{11} \Rightarrow a^2+\left(\dfrac{3a+37}{11}\right)^2+3a-14 = 0$ $\Rightarrow 2a^2+9a-5 = 0 \Rightarrow (a+5)(2a-1) = 0 \Rightarrow a = ...$	**dependent on both previous M marks.** Solves their equations simultaneously to obtain at least one value of $b = ...$ or $a = ...$	ddM1
	$z = \dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$ **only**	e.g. $\dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$ **or** $\dfrac{7}{2}\mathrm{i}+\dfrac{1}{2}$ **or** $0.5+3.5\mathrm{i}$	A1
			(5)

Question Number	Scheme	Notes	Marks
5.	$\dfrac{2z+3}{z+5-2\mathrm{i}} = 1+\mathrm{i}$		
(i) **Way 5**	$\dfrac{2z+3}{1+\mathrm{i}} = z+5-2\mathrm{i}$		
	$\dfrac{(2z+3)}{(1+\mathrm{i})}\dfrac{(1-\mathrm{i})}{(1-\mathrm{i})} = z+5-2\mathrm{i}$	Multiplies $\dfrac{(2z+3)}{(1+\mathrm{i})}$ by $\dfrac{(1-\mathrm{i})}{(1-\mathrm{i})}$ and sets equal to $z+5-2\mathrm{i}$	M1
	$\dfrac{(2z+3)(1-\mathrm{i})}{2} = z+5-2\mathrm{i}$		
	$2z+3-2\mathrm{i}z-3\mathrm{i} = 2z+10-4\mathrm{i}$		
	$2\mathrm{i}z = -7+\mathrm{i}$	**dependent on the previous M mark** Rearranges to make $2\mathrm{i}z = ...$	dM1
		Correct expression for $2\mathrm{i}z = ...$	A1
	$-2z = -7\mathrm{i}-1 \Rightarrow z = ...$	**dependent on both previous M marks** Multiplies both sides by i and attempts to find $z = ...$	ddM1
	$z = \dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$	e.g. $\dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$ **or** $\dfrac{7}{2}\mathrm{i}+\dfrac{1}{2}$ **or** $0.5+3.5\mathrm{i}$	A1
			(5)

	Question 5 Notes	
5. (i)	**Note**	Way 4 method generates $z = \dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$ and $z = -5+2\mathrm{i}$ but $z = \dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$ must be stated as the only answer for the final A mark
	Note	Give final A0 for a correct $a = \dfrac{1}{2}, b = \dfrac{7}{2}$ followed by an incorrect $\{z =\}\ \dfrac{7}{2}+\dfrac{1}{2}\mathrm{i}$
	Note	$\{z =\}\ \dfrac{1}{2}+\mathrm{i}\dfrac{7}{2}$ is fine for the final A mark
	Note	Give final A0 for $\{z =\}\ \dfrac{1+7\mathrm{i}}{2}$ without reference to e.g. $a = \dfrac{1}{2}, b = \dfrac{7}{2}$ **or** $\dfrac{1}{2}+\dfrac{7}{2}\mathrm{i}$, etc.
(ii)	**Note**	$w = (6-\lambda)+(3+2\lambda)\mathrm{i} \Rightarrow (15)^2 = (6-\lambda)^2-(3+2\lambda)^2$ is 1st M0
	Note	$\lvert(3+\lambda\mathrm{i})(2+\mathrm{i})\rvert = 15 \Rightarrow \sqrt{(3^2-\lambda^2)}\sqrt{(2^2-1^2)} = 15$ is 1st M0
	Note	Give final A0 for either • $\lambda = 6, -6 \Rightarrow \lambda = 6$ • $\lambda = 6, -6 \Rightarrow \lambda = -6$

Question Number	Scheme	Notes	Marks
6.	$C: y^2 = 32x$; S is the focus of C; $P(2, 8)$ lies on C; T lies on the direcrix of C. $H: xy = 4$		
(a)	S has coordinates $(8, 0)$	$(8, 0)$	B1 **cao**
			(1)
(b)	$\{PT$ is parallel to the x-axis $\Rightarrow\}$ $T(-8, 8) \Rightarrow PT = 2--8 = 10$ Focus-directrix Property $\Rightarrow PT = \sqrt{8^2 + (8-2)^2} = 10$	$PT = 10$	B1 **cao**
			(1)
(c)	$y = \sqrt{32}\,x^{\frac{1}{2}} \Rightarrow \frac{dy}{dx} = \frac{1}{2}\sqrt{32}\,x^{-\frac{1}{2}}$ or $2\sqrt{2}\,x^{-\frac{1}{2}}$	$\frac{dy}{dx} = \pm k x^{-\frac{1}{2}}; k \neq 0$	M1
	$y^2 = 32x \Rightarrow 2y\frac{dy}{dx} = 32$	$\lambda y \frac{dy}{dx} = \mu$; $\lambda, \mu \neq 0$	
	$x = 8t^2$, $y = 16t \Rightarrow \frac{dy}{dx} = \frac{dy}{dt}.\frac{dt}{dx} = 16\left(\frac{1}{16t}\right)$	$x = at^2$, $y = 2at \Rightarrow$ their $\frac{dy}{dt} \times \frac{1}{\text{their } \frac{dy}{dt}}$; $a \neq 0$	
	So at P, $m_T = 2$	Correct calculus work leading to $m_T = 2$	A1
	Either • $y - 8 = "2"(x-2)$ • $8 = "2"(2) + c \Rightarrow y = "2"x +$ their c	Correct straight line method using their gradient m_T $(\neq m_N)$ which is found by using calculus. **Note:** m_T must be a value	M1
	Correct algebra leading to $y = 2x+4$ *	Correct solution only	A1 *
			(4)
(d)	$x(2x+4) = 4$ or $\left(\frac{y-4}{2}\right)y = 4$ $\frac{4}{x} = 2x+4$ or $y = 2\left(\frac{4}{y}\right)+4$ $\frac{2}{t} = 2(2t)+4$	Subsitutes either • $y = 2x+4$ into $xy = 4$ • $y = \frac{4}{x}$ or $x = \frac{4}{y}$ into $y = 2x+4$ • $x = 2t$ and $y = \frac{2}{t}$ into $y = 2x+4$ to form an equation in either x only, y only or t only	M1
	$2x^2+4x-4=0$ or $x^2+2x-2=0$ or $\frac{1}{2}y^2-2y-4=0$ or $y^2-4y-8=0$ or $4t^2+4t-2=0$ or $2t^2+2t-1=0$	A correct 3 term quadratic **Note:** $2x^2+4x=4$, $\frac{1}{2}y^2-2y-4=0$, $2=4t^2+4t$ or $2x^2+4x-4\{=0\}$ are acceptable for this mark	A1
	• $\{x^2+2x-2=0 \Rightarrow\}$ $(x+1)^2-1-2=0 \Rightarrow x=...$ • $\{2t^2+2t-1=0 \Rightarrow\}$ $t = \frac{-2 \pm \sqrt{(2)^2-4(2)(-1)}}{2(2)}$ **and** either $x = 2\left(\frac{1}{2} \pm \frac{1}{2}\sqrt{3}\right)$ or $y = \frac{2}{\left(\frac{1}{2} \pm \frac{1}{2}\sqrt{3}\right)}$ • $\{y^2-4y-8=0 \Rightarrow\}$ $y = \frac{--4 \pm \sqrt{(-4)^2-4(1)(-8)}}{2(1)}$	**dependent on the previous M mark** Correct method (e.g. completing the square, applying the quadratic formula or factorising) of solving a 3TQ to find either $x = ...$ or $y = ...$	dM1
	Either $x = -1 \pm \sqrt{3}$ or $y = 2 \pm 2\sqrt{3}$	Both correct x coordinates or both correct y coordinates. **(See note)**	A1
	E.g. $y = 2(-1+\sqrt{3})+4$ or $y = \frac{4}{(-1+\sqrt{3})}$, etc.	**dependent on the first M mark** At least one attempt to find the other coordinate	dM1
	Either $(-1+\sqrt{3}, 2+2\sqrt{3}), (-1-\sqrt{3}, 2-2\sqrt{3})$ **or** $x = -1+\sqrt{3}$, $y = 2+2\sqrt{3}$ and $x = -1-\sqrt{3}$, $y = 2-2\sqrt{3}$	All correct and paired	A1
			(6)
			12

	Question 6 Notes	
6. (d)	**Note**	Condone $y = 2 \pm \sqrt{12}$ for the 2nd A1 mark.
	Note	Do not allow $(-1+\sqrt{3}, 2+\sqrt{12}), (-1-\sqrt{3}, 2-\sqrt{12})$ for the final A mark.
	Note	Writing $x = -1 \pm \sqrt{3}$, $y = 2 \pm 2\sqrt{3}$ without any evidence of the correct coordinate pairings is final A0
	Note	**<u>Writing coordinates the wrong way round</u>** E.g. writing $x = -1+\sqrt{3}$, $y = 2+2\sqrt{3}$ and $x = -1-\sqrt{3}$, $y = 2-2\sqrt{3}$ followed by $(-1+\sqrt{3}, 2-2\sqrt{3}), (-1-\sqrt{3}, 2+2\sqrt{3})$ is final A0
	Note	Imply the 1st dM1 mark for ***writing down*** the ***correct*** roots for ***their*** quadratic equation. E.g. • $2x^2 + 4x - 4 = 0$ or $x^2 + 2x - 2 = 0$ or $2x^2 + 4x = 4 \rightarrow x = -1 \pm \sqrt{3}$ • $\frac{1}{2}y^2 - 2y - 4 = 0$ or $y^2 - 4y - 8 = 0 \rightarrow y = 2 \pm 2\sqrt{3}$
	Note	You can imply the 1st A1, 1st dM1, 2nd A1 marks for either • $x(2x+4) = 4$ or $\frac{4}{x} = 2x + 4 \rightarrow x = -1 \pm \sqrt{3}$ • $\left(\frac{y-4}{2}\right)y = 4$ or $y = 2\left(\frac{4}{y}\right) + 4 \rightarrow y = 2 \pm 2\sqrt{3}$ with no intermediate working.
	Note	You can imply the 1st A1, 1st dM1, 2nd A1, 2nd dM1 marks for either • $x(2x+4) = 4$ or $\frac{4}{x} = 2x + 4 \rightarrow x = -1 \pm \sqrt{3}$ **and** $y = 2 \pm 2\sqrt{3}$ • $\left(\frac{y-4}{2}\right)y = 4$ or $y = 2\left(\frac{4}{y}\right) + 4 \rightarrow y = 2 \pm 2\sqrt{3}$ **and** $x = -1 \pm \sqrt{3}$ with no intermediate working. You can then imply the final A1 mark if they correctly state the correct coordinate pairings.
	Note	**2nd A1:** Allow this mark for both correct x coordinates or both correct y coordinates which are in the form $\frac{a \pm b\sqrt{c}}{d}$, where a, b, c and d are simplified integers

Question Number	Scheme	Notes	Marks
7.	$\mathbf{A}=\begin{pmatrix}6 & k\\ -3 & -4\end{pmatrix}, k\neq 8;\ \mathbf{A}^2+3\mathbf{A}^{-1}=\begin{pmatrix}5 & 9\\ -3 & -5\end{pmatrix};\ \mathbf{M}=\begin{pmatrix}-\frac{1}{2} & -\sqrt{3}\\ \frac{\sqrt{3}}{2} & -1\end{pmatrix}$		
(i)(a)	$\det(\mathbf{A})=6(-4)-(k)(-3)\ \ \{=-24+3k\}$	Correct $\det(\mathbf{A})$ which can be un-simplifed or simplifed	B1
	$\{\mathbf{A}^{-1}=\}\ \ \frac{1}{3k-24}\begin{pmatrix}-4 & -k\\ 3 & 6\end{pmatrix}$	$\begin{pmatrix}-4 & -k\\ 3 & 6\end{pmatrix}$	M1
		Correct $\mathbf{A}^{-1}$	A1
			(3)
(b)	$\{\mathbf{A}^2=\}\ \begin{pmatrix}36-3k & 6k-4k\\ -18+12 & -3k+16\end{pmatrix}\ \left\{=\begin{pmatrix}36-3k & 2k\\ -6 & -3k+16\end{pmatrix}\right\}$	Correct $\mathbf{A}^2$ which can be un-simplifed or simplifed	B1
			(1)
(c)	• $\begin{pmatrix}36-3k & 2k\\ -6 & -3k+16\end{pmatrix}+\frac{3}{3k-24}\begin{pmatrix}-4 & -k\\ 3 & 6\end{pmatrix}=\begin{pmatrix}5 & 9\\ -3 & -5\end{pmatrix}$ • $36-3k-\frac{12}{3k-24}=5$ • $2k-\frac{3k}{3k-24}=9$ • $-6+\frac{9}{3k-24}=-3$ • $-3k+16+\frac{18}{3k-24}=-5$ **Either** • attempts to form an equation for $(\text{their }\mathbf{A}^2)+3(\text{their }\mathbf{A}^{-1})=\begin{pmatrix}5 & 9\\ -3 & -5\end{pmatrix}$ in k • **or** attempts to add an element of $(\text{their }\mathbf{A}^2)$ to the corresponding element of $3(\text{their }\mathbf{A}^{-1})$ and equates to the corresponding element of the given matrix to form an equation in k		M1
	$\left\{\text{e.g. } -6+\frac{9}{3k-24}=-3\right\}\Rightarrow k=9$	**dependent on the previous M mark** Solves their equation to give $k=...$	dM1
		Final answer of $k=9$ ***only***	A1
			(3)
	Note: Parts (ii)(a) and (ii)(b) can be marked together **Please refer to the notes on the next page when marking (ii)(a) and (ii)(b)**		
(ii)(a)	• $p=\left(-\frac{1}{2}\right)(-1)-\left(-\sqrt{3}\right)\left(\frac{\sqrt{3}}{2}\right)=2$ • $-p\sin\theta=-\sqrt{3}$, $p\cos\theta=-1$ o $p=\sqrt{(\pm\sqrt{3})^2+(-1)^2}=2$ o $p=\frac{-\sqrt{3}}{-\sin"120°"}=2$ or $p=\frac{-1}{\cos"120°"}=2$	Attempts $p=\pm\frac{1}{2}\pm\left(\sqrt{3}\right)\left(\frac{\sqrt{3}}{2}\right)$ **or** uses a full method of trigonometry to find $p=...$	M1
		$p=2$ only	A1
			(2)
(b)	$\cos\theta=-\frac{1}{2},\ \sin\theta=\frac{\sqrt{3}}{2},\ \tan\theta=-\sqrt{3}$ E.g. • $\Rightarrow\theta=120°$ • $\Rightarrow\theta=180-\sin^{-1}\left(\frac{\sqrt{3}}{2}\right)=120°$ • $\Rightarrow\theta=180-\tan^{-1}\left(\sqrt{3}\right)=120°$	Uses trigonometry to find an expression or value for θ which is in the range $(1.57...,\ 3.14...)$ or $(90°,\ 180°)$ $(-3.14...,\ -4.71...)$ or $(-180°,\ -270°)$	M1
		$120°$ or $-240°$ or $\frac{2\pi}{3}$ or $-\frac{4\pi}{3}$ or awrt 2.09 or awrt -4.19	A1
			(2)
			11

	Question 7 Notes	
7. (i)(c)	**Note**	Give 1st M1 for $\begin{pmatrix} 36-3k-\dfrac{12}{3k-24} & 2k-\dfrac{3k}{3k-24} \\ -6+\dfrac{9}{3k-24} & -3k+16-\dfrac{18}{3k-24} \end{pmatrix} = \begin{pmatrix} 5 & 9 \\ -3 & -5 \end{pmatrix}$
	Note	• $36-3k-\dfrac{12}{3k-24}=5 \rightarrow 3k^2-55k+252=0 \rightarrow (k-9)(3k-28)=0 \rightarrow k=9, \dfrac{28}{3}$ • $2k-\dfrac{3k}{3k-24}=9 \rightarrow k^2-13k+36=0 \rightarrow (k-9)(k-4)=0 \rightarrow k=9, 4$ • $-6+\dfrac{9}{3k-24}=-3 \rightarrow k=9$ • $-3k+16-\dfrac{18}{3k-24}=-5 \rightarrow k^2-15k+54=0 \rightarrow (k-9)(k-6)=0 \rightarrow k=9, 6$
	Note	Uses a correct element equation in part (c) leading to $k=9$ is M1 dM1 A1 even if they have followed through an incorrect $\mathbf{A}^{-1}$ in (i)(a) or an incorrect $\mathbf{A}^2$ in (ii)(b).
	Note	Give M0 dM0 A0 for an incorrect method of $36-3k-4=5 \Rightarrow k=9$
(ii)	**Note**	$\mathbf{M}=\begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix}\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}=\begin{pmatrix} -\frac{1}{2} & -\sqrt{3} \\ \frac{\sqrt{3}}{2} & -1 \end{pmatrix} \Rightarrow \begin{pmatrix} \cos\theta & -p\sin\theta \\ \sin\theta & p\cos\theta \end{pmatrix}=\begin{pmatrix} -\frac{1}{2} & -\sqrt{3} \\ \frac{\sqrt{3}}{2} & -1 \end{pmatrix}$
	Note	**IMPORTANT NOTE** Give (ii)(a) M0A0 (b) M0A0 for a method of $\mathbf{M}=\begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}\begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix}=\begin{pmatrix} -\frac{1}{2} & -\sqrt{3} \\ \frac{\sqrt{3}}{2} & -1 \end{pmatrix} \Rightarrow \begin{pmatrix} \cos\theta & -\sin\theta \\ p\sin\theta & p\cos\theta \end{pmatrix}=\begin{pmatrix} -\frac{1}{2} & -\sqrt{3} \\ \frac{\sqrt{3}}{2} & -1 \end{pmatrix}$ leading to (ii)(a) $p=...$, (ii)(b) $\theta=...$
(ii)(a)	**Note**	$\det(\mathbf{M})=\left(-\dfrac{1}{2}\right)(-1)-\left(-\sqrt{3}\right)\left(\dfrac{\sqrt{3}}{2}\right)=2$ followed by $p=\sqrt{2}$ is M0 A0
	Note	$p=\det(\mathbf{M})=\left(-\dfrac{1}{2}\right)(-1)-\left(-\sqrt{3}\right)\left(\dfrac{\sqrt{3}}{2}\right)=2$ is M1 A1
	Note	$p=\dfrac{\sqrt{(\pm\sqrt{3})^2+(-1)^2}}{\sqrt{\left(-\dfrac{1}{2}\right)^2+\left(\dfrac{\sqrt{3}}{2}\right)^2}}=2$ is M1 A1

Question Number	Scheme	Notes	Marks
8.	(i) $u_1=3,\ u_{n+1}=u_n+3n-2$, $u_n=\frac{3}{2}n^2-\frac{7}{2}n+5$	(ii) $f(n)=3^{2n+3}+40n-27$ is divisible by 64	
(i)	$n=1,\ u_1=\frac{3}{2}-\frac{7}{2}+5=3$	Uses $u_n=\frac{3}{2}n^2-\frac{7}{2}n+5$ to show that $u_1=3$	B1
	(Assume the result is true for $n=k$)		
	$\{u_{k+1}=u_k+3k-2\Rightarrow\}$ $u_{k+1}=\frac{3}{2}k^2-\frac{7}{2}k+5+3k-2\ \left\{=\frac{3}{2}k^2-\frac{1}{2}k+3\right\}$	Finds u_{k+1} by attempting to substitute $u_k=\frac{3}{2}k^2-\frac{7}{2}k+5$ into $u_{k+1}=u_k+3k-2$. Condone one slip.	M1
	$=\frac{3}{2}(k+1)^2-3k-\frac{3}{2}-\frac{1}{2}k+3$	**dependent on the previous M mark**. Attempts to write u_{k+1} in terms of $(k+1)$	dM1
	$=\frac{3}{2}(k+1)^2-\frac{7}{2}k+\frac{3}{2}$		
	$=\frac{3}{2}(k+1)^2-\frac{7}{2}(k+1)+5$	***Uses algebra*** to achieve this result with no errors	A1
	If the result is true for $n=k$, then it is true for $n=k+1$. As the result has been shown to be true for $n=1$, then the result is true for all n $(\in\mathbb{Z}^+)$		A1 **cso**
			(5)
(ii) **Way 1**	$f(1)=3^5+40-27=256$	$f(1)=256$ is the minimum	B1
	$f(k+1)-f(k)=(3^{2(k+1)+3}+40(k+1)-27)-(3^{2k+3}+40k-27)$	Attempts $f(k+1)$ - $f(k)$	M1
	$f(k+1)-f(k)=8(3^{2k+3})+40$		
	$=8(3^{2k+3}+40k-27)-64(5k-4)$	$8(3^{2k+3}+40k-27)$ or $8f(k)$	A1
	or $=8(3^{2k+3}+40k-27)-320k+256$	$-64(5k-4)$ or $-320k+256$	A1
	$f(k+1)=8f(k)-64(5k-4)+f(k)$ or $f(k+1)=8f(k)-320k+256+f(k)$ or $f(k+1)=9(3^{2k+3}+40k-27)-320k+256$	**dependent on at least one of the previous accuracy marks being awarded.** Makes $f(k+1)$ the subject and expresses it in terms of $f(k)$ or $(3^{2k+3}+40k-27)$	dM1
	If the result is true for $n=k$, then it is true for $n=k+1$, As the result has been shown to be true for $n=1$, then the result is true for all n $(\in\mathbb{Z}^+)$		A1 **cso**
			(6)
(ii) **Way 2**	$f(1)=3^5+40-27=256$	$f(1)=256$ is the minimum	B1
	$f(k+1)=3^{2(k+1)+3}+40(k+1)-27$	Attempts $f(k+1)$	M1
	$f(k+1)=9(3^{2k+3})+40k+13$		
	$=9(3^{2k+3}+40k-27)-64(5k-4)$	$9(3^{2k+3}+40k-27)$ or $9f(k)$	A1
	or $=9(3^{2k+3}+40k-27)-320k+256$	$-64(5k-4)$ or $-320k+256$	A1
	$f(k+1)=9f(k)-64(5k-4)$ or $f(k+1)=9f(k)-320k+256$ or $f(k+1)=9(3^{2k+3}+40k-27)-320k+256$	**dependent on at least one of the previous accuracy marks being awarded.** Makes $f(k+1)$ the subject and expresses it in terms of $f(k)$ or $(3^{2k+3}+40k-27)$	dM1
	If the result is true for $n=k$, then it is true for $n=k+1$, As the result has been shown to be true for $n=1$, then the result is true for all n $(\in\mathbb{Z}^+)$		A1 **cso**
			11

Question Number	Scheme	Notes	Marks
8.	(ii) $f(n) = 3^{2n+3} + 40n - 27$ is divisible by 64		
(ii) **Way 3**	**General Method:** Using $f(k+1)$ - $mf(k)$; where m is an integer		
	$f(1) = 3^5 + 40 - 27 = 256$	$f(1) = 256$ is the minimum	B1
	$f(k+1) - mf(k) = (3^{2(k+1)+3} + 40(k+1) - 27) - m(3^{2k+3} + 40k - 27)$	Attempts $f(k+1)$ - $mf(k)$	M1
	$f(k+1) - mf(k) = (9-m)(3^{2k+3}) + 40k(1-m) + (13+27m)$		
	$= (9-m)(3^{2k+3} + 40k - 27) - 64(5k-4)$	$(9-m)(3^{2k+3} + 40k - 27)$ or $(9-m)f(k)$	A1
	or $= (9-m)(3^{2k+3} + 40k - 27) - 320k + 256$	$-64(5k-4)$ or $-320k+256$	A1
	$f(k+1) = (9-m)f(k) - 64(5k-4) + mf(k)$ or $f(k+1) = (9-m)f(k) - 320k + 256 + mf(k)$	**dependent on at least one of the previous accuracy marks being awarded.** Makes $f(k+1)$ the subject and expresses it in terms of $f(k)$ or $(3^{2k+3} + 40k - 27)$	dM1
	If the result is true for $n = k$, then it is true for $n = k+1$, As the result has been shown to be true for $n = 1$, then the result is is true for all n $(\in \mathbb{Z}^+)$		A1 **cso**
(ii) **Way 4**	**General Method:** Using $f(k+1)$ - $mf(k)$		
	$f(1) = 3^5 + 40 - 27 = 256$	$f(1) = 256$ is the minimum	B1
	$f(k+1) - mf(k) = (3^{2(k+1)+3} + 40(k+1) - 27) - m(3^{2k+3} + 40k - 27)$	Attempts $f(k+1)$ - $mf(k)$	M1
	$f(k+1) - mf(k) = (9-m)(3^{2k+3}) + 40k(1-m) + (13+27m)$		
	$m = -55 \Rightarrow f(k+1) + 55f(k) = 64(3^{2k+3}) - 2240k + 1472$	$m = -55$ and $64(3^{2k+3})$	A1
		$m = -55$ and $-2240k + 1472$	A1
	$f(k+1) = 64(3^{2k+3}) - 2240k + 1472 - 55f(k)$ or $f(k+1) = 64(3^{2k+3}) - 64(35k-23) - 55f(k)$	**dependent on at least one of the previous accuracy marks being awarded.** Makes $f(k+1)$ the subject and expresses it in terms of $f(k)$	dM1
	If the result is true for $n = k$, then it is true for $n = k+1$, As the result has been shown to be true for $n = 1$, then the result is true for all n $(\in \mathbb{Z}^+)$		A1 **cso**

Question 8 Notes

(i) & (ii)	**Note**	**Final A1 for parts (i) and (ii)** is dependent on all previous marks being scored in that part. It is gained by candidates conveying the ideas of **all** four underlined points **either** at the end of their solution **or** as a narrative in their solution.
(i)	**Note**	Moving from either $u_{k+1} = \frac{3}{2}k^2 - \frac{7}{2}k + 5 + 3k - 2$ or $u_{k+1} = \frac{3}{2}k^2 - \frac{1}{2}k + 3$ to $u_{k+1} = \frac{3}{2}(k+1)^2 - \frac{7}{2}(k+1) + 5$ ***with no intermediate stage involving either*** • writing u_{k+1} as a function of $(k+1)$ • **or** writing u_{k+1} as $u_{k+1} = \frac{3}{2}k^2 + 3k + \frac{3}{2} - \frac{7}{2}k - \frac{7}{2} + 5$ is dM1A0A0
	Note	Some candidates will write down $u_{k+1} = \frac{3}{2}k^2 - \frac{7}{2}k + 5 + 3k - 2$ **(give 1st M1)** and simplify this to $u_{k+1} = \frac{3}{2}k^2 - \frac{1}{2}k + 3$ They will then write $u_{k+1} = \frac{3}{2}(k+1)^2 - \frac{7}{2}(k+1) + 5$ **(give 2nd M1)** and use algebra to show that $u_{k+1} = \frac{3}{2}(k+1)^2 - \frac{7}{2}(k+1) + 5 = \frac{3}{2}(k^2 + 2k + 1) - \frac{7}{2}k - \frac{7}{2} + 5 = \frac{3}{2}k^2 - \frac{1}{2}k + 3$ **(give 1st A1)**

	Question 8 Notes Continued	
8. (ii)	**Note**	Some candidates may set $f(k) = 64M$ and so may prove the following general result • $\{f(k+1) = 9f(k) - 64(5k-4)\} \Rightarrow f(k+1) = 576M - 64(5k-4)$ • $\{f(k+1) = 9f(k) - 320k + 256\} \Rightarrow f(k+1) = 576M - 320k + 256$
	Note	$f(n) = 3^{2n+3} + 40n - 27$ can be rewritten as either $f(n) = 27(3^{2n}) + 40n - 27$ or $f(n) = 27(9^n) + 40n - 27$
	Note	In part (ii), Way 4 there are many alternatives where candidates focus on isolating $\beta(3^{2k+3})$ where β is a multiple of 64. Listed below are some alternative results: • $f(k+1) = 128(3^{2k+3}) - 119f(k) + 4800k - 3200$ • $f(k+1) = -64(3^{2k+3}) + 73f(k) - 2880k + 1984$ See below for how these are derived.

8. (ii)	(ii) $f(n) = 3^{2n+3} + 40n - 27$ is divisible by 64		
	The A1A1dM1 marks for Alternatives using $f(k+1)$ - $mf(k)$		
Way 4.1	$f(k+1) = 9(3^{2k+3}) + 40k + 13$		
	$= 128(3^{2k+3}) - 119(3^{2k+3}) + 40k + 13$		
	$= 128(3^{2k+3}) - 119[3^{2k+3} + 40k - 27] + 4800k - 3200$	$m = -119$ and $128(3^{2k+3})$	A1
		$m = -119$ and $4800k - 3200$	A1
	$f(k+1) = 128(3^{2k+3}) - 119f(k) + 4800k - 3200$ or $f(k+1) = 128(3^{2k+3}) - 119[3^{2k+3} + 40k - 27] + 4800k - 3200$	as before	dM1
Way 4.2	$f(k+1) = 9(3^{2k+3}) + 40k + 13$		
	$= -64(3^{2k+3}) + 73(3^{2k+3}) + 40k + 13$		
	$= -64(3^{2k+3}) + 73[3^{2k+3} + 40k - 27] - 2880k + 1984$	$m = 73$ and $-64(3^{2k+3})$	A1
		$m = 73$ and $-2880k + 1984$	A1
	$f(k+1) = -64(3^{2k+3}) + 73f(k) - 2880k + 1984$ or $f(k+1) = -64(3^{2k+3}) + 73[3^{2k+3} + 40k - 27] - 2880k + 1984$	as before	dM1

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom