edexcel

# Mark Scheme (Results)

## Summer 2016

## Pearson Edexcel GCE in Further Pure Mathematics 1 (6667/01)

**Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

**Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2016
Publications Code 6667_01_1606_MS
All the material in this publication is copyright
© Pearson Education Ltd 2016

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

# PEARSON EDEXCEL GCE MATHEMATICS

## General Instructions for Marking

1. The total number of marks for the paper is 75

2. The Edexcel Mathematics mark schemes use the following types of marks:

- **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
- **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
- **B** marks are unconditional accuracy marks (independent of M marks)
- Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
- ft – follow through
- the symbol $\surd$ will be used for correct ft
- cao – correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- oe – or equivalent (and appropriate)
- d… or dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- $*$ The answer is printed on the paper or ag- answer given
- $\sqsubset$ or d… The second mark is dependent on gaining the first mark

4. All A marks are ‘correct answer only’ (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.

6. If a candidate makes more than one attempt at any question:
   - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
   - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.

7. Ignore wrong working or incorrect statements following a correct answer.

# General Principles for Further Pure Mathematics Marking

*(But note that specific mark schemes may sometimes override these general principles).*

## Method mark for solving 3 term quadratic:

### 1. Factorisation

$(x^2+bx+c)=(x+p)(x+q)$, where $|pq|=|c|$, leading to x = ...

$(ax^2+bx+c)=(mx+p)(nx+q)$, where $|pq|=|c|$ and $|mn|=|a|$ , leading to x = ...

### 2. Formula

Attempt to use the correct formula (with values for a, b and c).

### 3. Completing the square

Solving $x^2+bx+c=0$: $\left(x\pm\frac{b}{2}\right)^2 \pm q \pm c=0,\ \ q\neq 0$ , leading to x = ...

## Method marks for differentiation and integration:

### 1. Differentiation

Power of at least one term decreased by 1. ($x^n \to x^{n-1}$)

### 2. Integration

Power of at least one term increased by 1. ($x^n \to x^{n+1}$)

## Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

## Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

| Question Number | Scheme | Marks |
|---|---|---|
| 1. | Determinant of **A** = $(1-k)(1+k)-k^2=0$ | M1 |
| | $1-k+k-k^2-k^2(=0)$<br>$1-2k^2(=0)$<br>So $k=\frac{\pm\sqrt{2}}{2}$ | A1<br><br>A1<br>(3) |
| | | **3 marks** |

Notes

M1: for attempting $ad-bc=0$ with $'=0'$ seen or **used** later in the solution.

A1: Correct (unsimplified) expression on LHS or correct equation after brackets expanded.

A1: Accept $\pm\frac{\sqrt{2}}{2}$, $\pm\frac{1}{\sqrt{2}}$, $\pm\sqrt{\frac{1}{2}}$ , $\pm\sqrt{0.5}$ . Must have $\pm$ for mark.

| Question Number | Scheme | Marks |
|---|---|---|
| 2. | (a) $f'(x) = \frac{9}{2}x^{\frac{1}{2}} + \frac{25}{2}x^{-\frac{3}{2}}$ | M1 A1 (2) |
| | (b) f(12.5)= 0.5115… (at least 0.51…) and $f'(12.5) = 16.1927...$ (at least 16… seen) | B1, B1 |
| | $x_1 = 12.5 - \frac{f(12.5)}{f'(12.5)} = 12.5 - \frac{0.5115}{16.1927...} = 12.468$ | M1 A1 (4) |
| | | **6 marks** |

Notes

(a) M1: for attempting differentiation i.e. decrease a power by 1
A1 Accept equivalent expression i.e. condone equivalent fractions.

(b) B1: One correct, must be explicitly seen if final answer incorrect, may be implied by correct final answer.
B1: Both correct; must be explicitly seen if final answer incorrect, may be implied by correct final answer.
M1:for attempting Newton- Raphson with their values for f(12.5) and $f'(12.5)$
A1: cao correct to 3dp
Newton Raphson used more than once – isw.

| Question Number | Scheme | Marks |
|---|---|---|
| 3. | (a) $\sum_{r=1}^{3n} r^2 = \frac{1}{6}3n(3n+1)(6n+1)$ or $\sum_{r=1}^{3n} r^2 = \frac{1}{2}n(3n+1)(6n+1)$ or equivalent | B1 (1) |
| | (b) See $\sum_{r=1}^{2n} r^2 = \frac{1}{3}n(2n+1)(4n+1)$ or equivalent | B1 |
| | Attempt to use $\sum_{r=1}^{3n} r^2 - \sum_{r=1}^{2n} r^2 = \frac{n}{6}\{3(3n+1)(6n+1) - 2(2n+1)(4n+1)\}$ | M1 |
| | $= \frac{n}{6}\{(54n^2+27n+3) - (16n^2+12n+2)\}$ | dM1 |
| | $= \frac{n}{6}\{(38n^2+15n+1)\}$ $(a=38, b=15, c=1)$ | A1 (4) |
| | | **5 marks** |

**Notes**

(a) B1: Either right hand side or exact equivalent - isw if expanded

(b) B1: States or uses $\sum_{r=1}^{2n} r^2 = \frac{1}{3}n(2n+1)(4n+1)$

M1: Subtracts their sum to $2n$ or $2n - 1$ **and** attempts to factorise by $\frac{n}{6}$ seen anywhere.

dM1: Expands two quadratics dependent on first M1

A1: cao

| Question Number | Scheme | Marks |
|---|---|---|
| 4. | (a) $z = \dfrac{4(1-\mathrm{i})}{(1+\mathrm{i})(1-\mathrm{i})}$ | M1 |
| | $z = 2(1-\mathrm{i})$ or $2 - 2\mathrm{i}$ or exact equivalent. | A1 (2) |
| | (b) $z^2 = (2-2\mathrm{i})(2-2\mathrm{i}) = 4 - 8\mathrm{i} + 4\mathrm{i}^2$ | M1 |
| | $= -8\mathrm{i}$ | A1 cao (2) |
| | (c) If $z$ is a root so is $z^*$ So $(x - 2 + 2\,\mathrm{i})(x - 2 - 2\,\mathrm{i})$ (or $x^2 - 2\,\mathrm{Re}(z).x + \lvert z\rvert^2$ **)** | M1 |
| | So $(x - 2 + 2\,\mathrm{i})(x - 2 - 2\,\mathrm{i}) = 0$ (or $x^2 - 2\,\mathrm{Re}(z).x + \lvert z\rvert^2 = 0$**)** and so $p = q =$ | M1 |
| | Equation is $x^2 - 4x + 8(= 0)$ or $p = -4$ and $q = 8$ | A1 (3) (7 marks) |
| ALT 1 | (c) | |
| | Substitutes $z = 2 - 2\mathrm{i}$ and $z^2 = -8\mathrm{i}$ into quadratic and equates real and imaginary parts to obtain $2p + q = 0$ and $-2p - 8 = 0$ | M1 |
| | Attempts to solve simultaneous equations to obtain $p = -4$ and $q = 8$ | M1A1 |
| ALT 2 | (c) | |
| | Attempts to obtain $p = -$ sum of roots | M1 |
| | Attempts product of roots to obtain $q =$ | M1 |
| | Equation is $x^2 - 4x + 8(= 0)$ or $p = -4$ and $q = 8$ | A1 |
| ALT 3 | (c) $x - 2 = \pm 2\mathrm{i}$ either sign acceptable | M1 |
| | $(x-2)^2 = -4 \Rightarrow x^2 - 4x + 4 = -4$ i.e square and attempt to expand to give 3-term quadratic | M1 |
| | Equation is $x^2 - 4x + 8(= 0)$ or $p = -4$ and $q = 8$ | A1 |

**Notes**

(a) M1: Multiplies numerator and denominator by $1 - \mathrm{i}$ or by $-1 + \mathrm{i}$
A1: cao

(b) M1: Squares their $z$, or the given $z = \dfrac{4}{1+\mathrm{i}}$, to produce at least 3 terms which can be implied by the correct answer.
A1: $-8\mathrm{i}$ or $0 - 8\mathrm{i}$ only

(c) M1: Uses their $z$ and $z^*$ in $(x - z)(x - z^*)$
M1: Multiplies two factors and obtains $p =$ or $q =$
A1: Both correct required – can be implied by $x^2 - 4x + 8$
ALT 1
(c) M1: Substitutes their $z$ and their $z^2$ into the quadratic and equates real and imaginary parts to obtain two equations in $p$ and $q$
M1: Attempts to solve for one unknown to obtain $p =$ or $q =$
A1: Both correct required – can be implied by $x^2 - 4x + 8(= 0)$

| Question Number | Scheme | Marks |
|---|---|---|
| 5. | (a) Gradient $= \frac{2ap-2aq}{ap^2-aq^2}$ seen | B1 |
| | $\left(\frac{2ap-2aq}{ap^2-aq^2}=\frac{2a(p-q)}{a(p-q)(p+q)}\right)=\frac{2}{p+q}$ or seen in an equation | B1 |
| | uses $y-y_1=m(x-x_1)$ to give $(y-2aq)="m"(x-aq^2)$ or equivalent with $p$<br>or uses $y = mx + c$ to give $y="m"x+c$ and substitute a point to find $c$ | M1 |
| | or uses $\frac{y-y_1}{y_2-y_1}=\frac{x-x_1}{x_2-x_1}$ to give $\frac{(y-2aq)}{2ap-2aq}=\frac{(x-aq^2)}{ap^2-aq^2}$ or equivalent with $p$ | |
| | so $(y-2aq)=\frac{2}{p+q}(x-aq^2)$ or $(y-2ap)=\frac{2}{p+q}(x-ap^2)$ or $y=\frac{2}{p+q}x+\frac{2apq}{p+q}$ or $\frac{(y-2aq)}{2a}=\frac{(x-aq^2)}{a(p+q)}$ | A1 |
| | See $2aq^2$ or $2ap^2$ term appear and disappear to give $y(p+q)=2x+2apq$ * | A1 cso (5) |
| | (b) Substitute ($a$ , 0 ) into line equation, to give $0=2a+2apq$ so $pq=-1$ | B1 (1) |
| | (c) $y=2a^{\frac{1}{2}}x^{\frac{1}{2}}\Rightarrow\frac{dy}{dx}=a^{\frac{1}{2}}x^{-\frac{1}{2}}$ or $y^2=4ax\Rightarrow 2y\frac{dy}{dx}=4a$ or $\frac{dy}{dx}=\frac{dy}{dp}\times\frac{dp}{dx}\Rightarrow\frac{dy}{dx}=2a\times\frac{1}{2ap}$ | M1 |
| | So at $P$ tangent gradient = $\frac{1}{p}$ | A1 (2) |
| | (d) At $Q$ tangent gradient = $\frac{1}{q}$ | B1 |
| | $\frac{1}{p}\times\frac{1}{q}=\frac{1}{pq}=\frac{1}{-1}=-1$ with at least one intermediate step, the tangents are **perpendicular** or at **right angles** | B1cso (2) (10 marks) |

Notes


(a) B1: Correct statement for gradient (isw) B1: $\frac{2}{p+q}$ - can be seen later in the solution.

M1: use of a correct formula for a line equation through $P$ or $Q$ with their gradient.
Must be finding a chord, not a tangent or a normal.
A1: for a correct line equation with simplified gradient in any equivalent form
A1: cso (as given answer)
(b) B1: For using ($a$, 0) to show that $pq$ = -1
(c) M1: Use calculus to find an expression for d$y$/d$x$ and substitute coordinates of $P$.
They may use chord gradient and let $p$ tend to $q$.
(d) B1: 1/$q$ seen B1: $\frac{1}{p}\times\frac{1}{q}=-1$ or $\frac{1}{p}=-\frac{1}{1/q}$ or $\frac{1}{q}=-\frac{1}{1/p}$ and at least words in bold with no errors seen.

| Question Number | Scheme | Marks |
|---|---|---|
| 6. | (a) **Rotation** ,135 degrees or $\frac{3\pi}{4}$ radians (anticlockwise) about $O$ or 225 degrees or $\frac{5\pi}{4}$ clockwise about $O$. | M1, A1 (2) |
| | (b) $\begin{pmatrix} -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix}\begin{pmatrix} p \\ q \end{pmatrix} = \begin{pmatrix} 6\sqrt{2} \\ 3\sqrt{2} \end{pmatrix}$ | |
| | $-p-q=12$ and $p-q=6$ or equivalent | M1A1 |
| | $p$ = –3 and $q$ = –9 or $\begin{pmatrix} -3 \\ -9 \end{pmatrix}$ | B1 cso (3) |
| | ALT Uses Inverse matrix $\mathbf{P}^{-1}$ with vector $= \begin{pmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix}\begin{pmatrix} 6\sqrt{2} \\ 3\sqrt{2} \end{pmatrix}$ | M1A1 |
| | $p$ = –3 and $q$ = –9 or $\begin{pmatrix} -3 \\ -9 \end{pmatrix}$ | B1 cso (3) |
| | (c) $\mathbf{Q} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ | B1 (1) |
| | Accept **T** if used instead of **R** | |
| | (d) $\mathbf{R} = "\mathbf{Q}"\mathbf{P} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}\begin{pmatrix} -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix}$ | M1 A1 A1 (3) |
| | (e) $\mathbf{R}^{-1} = \frac{1}{-1}\begin{pmatrix} -\frac{1}{\sqrt{2}} & +\frac{1}{\sqrt{2}} \\ +\frac{1}{\sqrt{2}} & +\frac{1}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix}$ | |
| | $= \mathbf{R}$ (so matrix is self inverse and so transformation is self inverse) | B1 **(1)** **10 marks** |
| | ALT 1 (e) $\mathbf{RR} = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix}\begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ **(**so **R** is self inverse and so transformation is self inverse) | B1 |
| | ALT 2 (e) Matrix represents a reflection (so is self inverse) | B1 |

**Notes**

(a) M1: Rotation only A1: 135 degrees about $O$
SC: 135 degrees about $O$ only award M1A0.
(b) M1: Multiplies matrices in correct order to obtain two equations in $p$ and $q$.
A1: Two correct equations
B1 cso: $p$ and $q$ both correct, may be in vector form. No errors seen in solution.
ALT (b) M1: Attempt to find Inverse Matrix and pre-multiply A1: Correct Inverse Matrix used
B1 cso: $p$ and $q$ both correct, may be in vector form. No errors seen in solution.
(d) M1: Sets matrix product correct way round and obtains one correct term for their **Q**
A1: Two correct terms from a correct **Q.** **Q** incorrect award A0 here. A1: Completely correct matrix
(e) B1: Calculates $\mathbf{R^{-1}} = \mathbf{R}$ or calculates $\mathbf{R}^2$ and indicates that $\mathbf{R}^2 = \mathbf{I}$ or states that **R** represents a reflection.

| Question Number | Scheme | Marks |
|---|---|---|
| 7. | (a) $z^2 = (a+2\text{i})(a+2\text{i}) = (a^2-4)+4\text{i}a$<br>So $z^2+2z = (a^2-4+2a)+\text{i}(4a+4)$ or $x = (a^2+2a-4)$ and $y = 4a+4$ | M1<br>M1 A1 A1<br>(4) |
| | (b) and so $4a+4=0 \rightarrow a=-1$ | B1<br>(1) |
| | ALT (b)<br>Substitute $a = -1$ and show that $y = 0$ | B1<br>B1 |
| | (c) $\lvert z\rvert = \sqrt{5}$ or awrt 2.24<br>$\arctan(-2) = 2.03$ | B1<br>M1, A1 cao<br>(3) |
| | (d)  | M1 A1<br>B1ft<br>(3) |
| | (e) OP and QR are parallel , and QR is twice the length of OP<br>Or Enlargement with Scale Factor 2 (centre O), followed by translation $\begin{pmatrix}-3\\-4\end{pmatrix}$<br>Or Enlargement with Scale Factor 2, centre (3,4) or centre 3 + 4i<br>$\overrightarrow{QR} = 2\overrightarrow{OP}$ with clear indication of vectors award B1B1, without vectors award B0B1 | B1, B1<br>(2)<br>**13 marks** |

**Notes:**

(a) M1: Squares $z$ to produce at least 3 terms which can be implied by the correct answer.
M1: Adds $2z$ to their $z^2$
A1: Correct $x$ A1 Correct $y$ accept $4ai+4i$
(b) B1: Completely accurate cao
(c) B1: $\sqrt{5}$ or 2.24 or awrt 2.24
M1 for using tan or arctan
A1 cao 2.03
(d) M1: **Either** their $OP$ in the correct quadrant labelled $P$ or $z$ or their $-1+2\text{i}$ or their (-1,2) or axes labelled **or** their $OQ$ in the correct quadrant labelled $Q$ or $z^2$ or their $-3-4\text{i}$ or their (-3, -4) or axes labelled
A1: Both $OP$ and $OQ$ correct i.e. in the 2nd and 3rd quadrants respectively.
B1ft: $OR: z^2+2z(=-5)$ **on real axis** to left of the origin.
Accept points or lines. Arrows not required. Axes need not be labelled Re and Im.
Treat correct quadrant (or on axis) as important aspect for accuracy, lengths of lines if present can be accepted as correct.

| Question Number | Scheme | Marks |
|---|---|---|
| 8. | (i) If $n=1$, $\sum_{r=1}^{n}\frac{2r+1}{r^2(r+1)^2}=\frac{3}{4}$ **and** $1-\frac{1}{(n+1)^2}=\frac{3}{4}$, so true for $n = 1$. | B1 |
| | Assume result true for $n = k$ **and** consider $\sum_{r=1}^{k+1}\frac{2r+1}{r^2(r+1)^2}=1-\frac{1}{(k+1)^2}+\frac{2(k+1)+1}{(k+1)^2(k+2)^2}$ | M1 |
| | $=1-\left(\frac{(k+2)^2}{(k+1)^2(k+2)^2}-\frac{2(k+1)+1}{(k+1)^2(k+2)^2}\right)=1-\left(\frac{(k^2+2k+1)}{(k+1)^2(k+2)^2}\right)$ | A1 |
| | $=1-\left(\frac{(k+1)^2}{(k+1)^2(k+2)^2}\right)=1-\left(\frac{1}{(k+1+1)^2}\right)$ | M1 |
| | **True** for $\boldsymbol{n = k + 1}$ if **true** for $\boldsymbol{n = k}$, ( and **true** for $\boldsymbol{n = 1}$) so **true** by induction for all $\boldsymbol{n \in Z^+}$ | A1cso (5) |
| | (ii) $n = 1$: $u_1=5\times\left(\frac{1}{3}\right)^1+\frac{4}{3}=3$ so expression for $u_n$ true for $n = 1$ | B1 |
| | Assume result true for $n = k$ **and** consider $u_{k+1}=\frac{1}{3}(5\times\left(\frac{1}{3}\right)^k+\frac{4}{3})+\frac{8}{9}$ | M1 |
| | Obtain $u_{k+1}=5\times\left(\frac{1}{3}\right)^{k+1}+\frac{4}{9}+\frac{8}{9}$ | A1 |
| | $5\times\left(\frac{1}{3}\right)^{k+1}+\frac{4}{3}$ and deduce that result is true for $n = k + 1$ | dM1 |
| | **True** for $\boldsymbol{n = k + 1}$ if **true** for $\boldsymbol{n = k}$, ( and **true** for $\boldsymbol{n = 1}$) so **true** by induction for all $\boldsymbol{n \in Z^+}$ | A1 cso (5) **10 marks** |

**Notes:**

(i) B1: Checks $n = 1$ on both sides **and** states true for $n = 1$ seen anywhere

M1: (Assumes true for) $n = k$ **and** adds $(k+1)^{th}$ term to sum of $k$ terms

A1: $1-\left(\frac{(k^2+2k+1)}{(k+1)^2(k+2)^2}\right)$ seen (linked to 2nd M)

M1: $(k+1)^2(k+2)^2$ attempted as common denominator of two fractions.

A1cso: Makes correct complete induction statement including at least statements in bold. Accept $n\geq 1$ or $n=1,2,3...$ or all positive Integers or all $n$. Statement true for $n = 1$ here could contribute to B1 mark earlier.

(ii) B1: Checks $n = 1$ in $u_n$ and states true for $n = 1$ seen anywhere.

M1: (Assumes result for) $n = k$ **and** substitutes $u_k$ into correct expression for $u_{k+1}$

A1: $\frac{4}{9}+\frac{8}{9}$ or $\frac{1}{3}.\frac{4}{3}+\frac{8}{9}$ seen

dM1: Obtains $5\times\left(\frac{1}{3}\right)^{k+1}+\frac{4}{3}$ and statement true for $n = k + 1$ or equivalent seen anywhere dependent on previous M.

A1cso: Makes correct complete induction statement including at least statements in bold. Accept $n\geq 1$ or $n=1,2,3...$ or all positive Integers or all $n$. Statement true for $n = 1$ here could contribute to B1 mark earlier.

| Question Number | Scheme | Marks |
|---|---|---|
| 9 (a) | $y=\frac{25}{x}\Rightarrow \frac{dy}{dx}=-25x^{-2}$ , | B1 |
| | **or** $y+x\frac{dy}{dx}=0\Rightarrow \frac{dy}{dx}=-\frac{y}{x}$ **or** $\dot{x}=5,\ \dot{y}=-\frac{5}{t^2}$ so $\frac{dy}{dx}=-\frac{1}{t^2}$ | |
| | and at $P$ $\frac{dy}{dx}=-\frac{1}{p^2}$ so gradient of normal is $p^2$ | M1 A1 |
| | **Either** $y-\frac{5}{p}=p^2(x-5p)$ **or** $y=p^2x+k$ and use $x=5p,\ y=\frac{5}{p}$ | M1 |
| | $\Rightarrow \quad y-p^2x=\frac{5}{p}-5p^3$ (*) | A1cso (5) |
| (b) | At the point $A$: $y+p^2y=\frac{5}{p}-5p^3$ or $-x-p^2x=\frac{5}{p}-5p^3$ | |
| | $y(1+p^2)=\frac{5}{p}(1-p^4)$ or $-x(1+p^2)=\frac{5}{p}(1-p^4)$ | M1 |
| | $y=\frac{\frac{5}{p}(1-p^2)(1+p^2)}{(1+p^2)}=\frac{5}{p}(1-p^2)=\frac{5}{p}-5p$ or $x=\frac{-\frac{5}{p}(1-p^2)(1+p^2)}{(1+p^2)}=\frac{-5}{p}(1-p^2)=\frac{-5}{p}+5p$ * | M1 |
| | so $x=-\frac{5}{p}(1-p^2)=-\frac{5}{p}+5p$ and $y=\frac{5}{p}-5p$ * | A1cso (3) |
| (c) | $M$ has coordinates $\left(-\frac{5}{2p}+5p,\ \frac{5}{p}-\frac{5p}{2}\right)$ o.e. | B1 |
| | So when $y=0$, $\frac{5}{p}-\frac{5p}{2}=0$ and $p=\sqrt{2}$ so $M$ has $x$ coordinate $\frac{15}{4}\sqrt{2}$ o.e. | M1 A1 (3) |
| | | **11 marks** |

Notes

(a) B1: Any correct expression for gradient of tangent
M1: Substitutes values into **derived expression using calculus** to give gradient of normal at $P$
A1: cao. Can be implied by use in equation of a straight line
M1: Use of formula for the equation of a straight line with their changed gradient
A1: cso
(b) M1: Replaces $x$ by $-y$ or $y$ by $-x$
M1: Factorises $(1-p^4)$ to simplify answer in first variable
A1 cso: Obtains both $x$ and $y$
ALT (b) Accept Verification.
M1: Substitutes the coordinates of $A$ into the equation of the normal
M1: Substitutes the coordinates of $A$ into both the normal and $y=-x$.
A1 cso: No errors seen
(c) B1: Correct $x$- coordinate of midpoint (may be implied) and correct $y$ coordinate, accept equivalent forms
M1: Puts their $y=0$ and finds value for $p$ to use in $x=$
A1: $+\frac{15}{4}\sqrt{2}$ or equivalent only

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London WC2R 0RL