GradeMax

edexcel

Mark Scheme (Results)

January 2014

Pearson Edexcel International Advanced Level

Further Pure Mathematics 1 (6667A/01)

GradeMax

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2014
Publications Code IA037751
All the material in this publication is copyright
© Pearson Education Ltd 2014

GradeMax

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

GradeMax

EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.

2. The Edexcel Mathematics mark schemes use the following types of marks:

- **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
- **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
- **B** marks are unconditional accuracy marks (independent of M marks)
- Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
- ft – follow through
- the symbol $\surd$ will be used for correct ft
- cao – correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- oe – or equivalent (and appropriate)
- dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- $*$ The answer is printed on the paper
- ⌐ The second mark is dependent on gaining the first mark

4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.

6. If a candidate makes more than one attempt at any question:
   - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
   - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.

7. Ignore wrong working or incorrect statements following a correct answer.

GradeMax

General Principles for Pure Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

Method mark for solving 3 term quadratic:

1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$, where $|pq| = |c|$, leading to x =

$(ax^2 + bx + c) = (mx + p)(nx + q)$, where $|pq| = |c|$ and $|mn| = |a|$, leading to x =

2. Formula

Attempt to use correct formula (with values for *a*, *b* and *c*).

3. Completing the square

Solving $x^2 + bx + c = 0$: $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c, \quad q \neq 0,$ leading to x =...

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \to x^{n-1}$)

2. Integration

Power of at least one term increased by 1. ($x^n \to x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.
Normal marking procedure is as follows:
Method mark for quoting a correct formula and attempting to use it, even if there are mistakes in the substitution of values.
Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required.

Question Number	Scheme	Notes	Marks
1.	$f(x) = 2x - 5\cos x$, x measured in radians		
(a)	$f(1) = -0.7015115293...$ $f(1.4) = 1.950164285...$	Either any one of $f(1) =$ awrt -0.7 or $f(1.4) = 1.9$ or awrt 2.0	M1
	Sign change (and $f(x)$ is continuous) therefore a root α exists between $x = 1$ and $x = 1.4$	both values correct, sign change and conclusion	A1 **[2]**
(b)	$f(1.2) = 0.5882112276...$ $\{\Rightarrow 1 \le \alpha \le 1.2\}$	$f(1.2) =$ awrt 0.6	B1
		Attempt to find $f(1.1)$	M1
	$f(1.1) = -0.06798060713...$ $\Rightarrow 1.1 \le \alpha \le 1.2$	$f(1.1) = -0.06$ or awrt -0.07 with $1.1 \le \alpha \le 1.2$ or $1.1 < \alpha < 1.2$ or $[1.1, 1.2]$ or $(1.1, 1.2)$.	A1 **[3]** **5**

Question Number	Scheme	Notes	Marks
2.	$\mathbf{A} = \begin{pmatrix} -4 & 10 \\ -3 & k \end{pmatrix}$, where k is a constant		
(i)	$\det \mathbf{A} = (-4)(k) - (-3)(10)$	Applies "$ad \pm bc$" to $\mathbf{A}$	M1
	$\Rightarrow -4k + 30 = 2$ or $-4k + 30 = -2$	Equates their $\det \mathbf{A}$ to either 2 or -2	dM1
		Either $k = 8$ or $k = 7$	A1
	$\Rightarrow k = 7$ or $k = 8$	Both $k = 8$ and $k = 7$	A1
			[4]
(ii)	$\mathbf{B} = \begin{pmatrix} 1 & -2 & 3 \\ -2 & 5 & 1 \end{pmatrix}$, $\mathbf{C} = \begin{pmatrix} 2 & 8 \\ 0 & 2 \\ 1 & -2 \end{pmatrix}$		
	$\mathbf{BC} = \begin{pmatrix} 1 & -2 & 3 \\ -2 & 5 & 1 \end{pmatrix}\begin{pmatrix} 2 & 8 \\ 0 & 2 \\ 1 & -2 \end{pmatrix} = \begin{pmatrix} 5 & -2 \\ -3 & -8 \end{pmatrix}$	Writes down a complete 2×2 matrix.	M1
		Any 3 out of 4 elements correct	A1
		Correct answer.	A1
			[3]
			7

Question Number	Scheme	Notes	Marks
3.	$x = 2t,\ y = \frac{2}{t},\ t \neq 0$		
	$t = \frac{1}{2} \Rightarrow P\left(1, 4\right),\quad t = 4 \Rightarrow Q\left(8, \frac{1}{2}\right)$	Coordinates for either P or Q are correctly stated. (Can be implied).	B1
	$m(PQ) = \frac{\frac{1}{2} - 4}{8 - 1}\ \left\{= -\frac{1}{2}\right\}$	An attempt to find the gradient of the chord PQ.	M1
	$m(L) = 2$	Applying $m(L) = \frac{-1}{\text{their } m(PQ)}$	M1
	So, $L : y = 2x$	$y = 2x$	A1 oe
			[4]
			4

Question Number	Scheme	Notes	Marks
4.	$f(x) = 2\sqrt{x} - \frac{6}{x^2} - 3, \quad x > 0$		
	$f'(x) = x^{-\frac{1}{2}} + 12x^{-3} \{+ 0\}$	$\pm\lambda x^{-\frac{1}{2}}$ or $\pm\mu x^{-3}$ Correct differentiation	M1 A1
	$f(3.5) = 0.2518614684...$ $\{f'(3.5) = 0.8144058657...\}$	f (3.5) = awrt 0.25	B1
	$\beta = 3.5 - \left(\frac{"0.2518614684..."}{"0.8144058657..."}\right)$ $= 3.190742075...$	Correct application of Newton-Raphson using their values.	M1
	$= 3.191$ (3dp)	3.191	A1 **cao** **[5]** **5**

Question Number	Scheme	Notes	Marks
5.	$z = 5 + \mathrm{i}\sqrt{3}, \quad w = \sqrt{3} - \mathrm{i}$		
(a)	$\lvert w\rvert = \left\{\sqrt{\left(\sqrt{3}\right)^2 + (-1)^2}\right\} = 2$	2	B1
			[1]
(b)	$zw = \left(5 + \mathrm{i}\sqrt{3}\right)\left(\sqrt{3} - \mathrm{i}\right)$		
	$= 5\sqrt{3} - 5\mathrm{i} + 3\mathrm{i} + \sqrt{3}$		
	$= 6\sqrt{3} - 2\mathrm{i}$	Either the real or imaginary part is correct.	M1
		$6\sqrt{3} - 2\mathrm{i}$	A1
			[2]
(c)	$\frac{z}{w} = \frac{\left(5 + \mathrm{i}\sqrt{3}\right)}{\left(\sqrt{3} - \mathrm{i}\right)} \times \frac{\left(\sqrt{3} + \mathrm{i}\right)}{\left(\sqrt{3} + \mathrm{i}\right)}$	Multiplies by $\frac{\left(\sqrt{3} + \mathrm{i}\right)}{\left(\sqrt{3} + \mathrm{i}\right)}$	M1
	$= \frac{5\sqrt{3} + 5\mathrm{i} + 3\mathrm{i} - \sqrt{3}}{3 + 1}$	Simplifies realising that a real number is needed on the denominator and applies $\mathrm{i}^2 = -1$ on their numerator expression and denominator expression.	M1
	$\left\{= \frac{4\sqrt{3} + 8\mathrm{i}}{4}\right\} = \sqrt{3} + 2\mathrm{i}$	$\sqrt{3} + 2\mathrm{i}$	A1
			[3]
(d)	$z + \lambda = 5 + \mathrm{i}\sqrt{3} + \lambda = (5 + \lambda) + \mathrm{i}\sqrt{3}$		
	$\left\{\arg(z + \lambda) = \frac{\pi}{3} \Rightarrow\right\} \quad \frac{\sqrt{3}}{5 + \lambda} = \tan\left(\frac{\pi}{3}\right)$	$\frac{\sqrt{3}}{\text{their combined real part}} = \tan\left(\frac{\pi}{3}\right)$	M1 oe
	$\left\{\frac{\sqrt{3}}{5 + \lambda} = \frac{\sqrt{3}}{1} \Rightarrow 5 + \lambda = 1 \Rightarrow\right\} \lambda = -4$	-4	A1
			[2]
			8

GradeMax

Question Number	Scheme	Notes	Marks
6. (a)	$\sum_{r=1}^{n} r(r+1)(r-1) = \sum_{r=1}^{n} (r^3 - r)$		
	$= \frac{1}{4}n^2(n+1)^2 - \frac{1}{2}n(n+1)$	An attempt to use at least one of the standard formulae correctly. Correct expression.	M1 A1
	$= \frac{1}{4}n(n+1)\left(n(n+1) - 2\right)$	An attempt to factorise out at least $n(n+1)$.	M1
	$= \frac{1}{4}n(n+1)\left(n^2 + n - 2\right)$		
	$= \frac{1}{4}n(n+1)(n-1)(n+2)$	Achieves the correct answer. (Note: $a = 2$).	A1
			[4]
(b)	$\sum_{r=1}^{n} r(r+1)(r-1) = 10\sum_{r=1}^{n} r^2$		
	$\frac{1}{4}n(n+1)(n-1)(n+2) = \frac{10}{6}n(n+1)(2n+1)$	Sets their part (a) $= \frac{10}{6}n(n+1)(2n+1)$	M1
	$\frac{1}{4}(n-1)(n+2) = \frac{5}{3}(2n+1)$		
	$3(n^2 + n - 2) = 20(2n+1)$		
	$3n^2 - 37n - 26 = 0$	Manipulates to a "3TQ = 0". $3n^2 - 37n - 26 = 0$	M1 A1
	$(3n + 2)(n - 13) = 0$	A valid method for factorising a 3TQ.	M1
	$n = 13$	Only one solution of $n = 13$	A1
			[5]
			9

Question Number	Scheme	Notes	Marks
7.	$\mathbf{P} = \begin{pmatrix} 3a & -2a \\ -b & 2b \end{pmatrix}$, $\mathbf{M} = \begin{pmatrix} -6a & 7a \\ 2b & -b \end{pmatrix}$		
(a)	$\mathbf{P}^{-1} = \frac{1}{4ab}; \begin{pmatrix} 2b & 2a \\ b & 3a \end{pmatrix}$	$\frac{1}{4ab}$ Two out of four elements correct. Correct matrix.	B1; M1 A1 **[3]**
(b)	$\mathbf{M} = \mathbf{PQ}$ $\Rightarrow \mathbf{P^{-1}M} = \mathbf{P^{-1}PQ} \Rightarrow \mathbf{Q} = \mathbf{P^{-1}M}$ $\mathbf{Q} = \frac{1}{4ab}\begin{pmatrix} 2b & 2a \\ b & 3a \end{pmatrix}\begin{pmatrix} -6a & 7a \\ 2b & -b \end{pmatrix}$ $= \frac{1}{4ab}\begin{pmatrix} -8ab & 12ab \\ 0 & 4ab \end{pmatrix}$ $= \begin{pmatrix} -2 & 3 \\ 0 & 1 \end{pmatrix}$	Multiples their $\mathbf{P^{-1}}$ by $\mathbf{M}$ Two out of four elements correct. Correct matrix.	M1 A1 A1 **[3]**
			6

Question Number	Scheme	Notes	Marks
8.	$y^2 = 4ax$, at $P\left(ap^2, 2ap\right)$.		
(a)	$y = 2\sqrt{a}\,x^{\frac{1}{2}} \Rightarrow \dfrac{dy}{dx} = \sqrt{a}\,x^{-\frac{1}{2}}$	$\dfrac{dy}{dx} = \pm k x^{-\frac{1}{2}}$	
	or (implicitly) $2y\dfrac{dy}{dx} = 4a$	or $ky\dfrac{dy}{dx} = c$	M1
	or (chain rule) $\dfrac{dy}{dx} = 2a \times \dfrac{1}{2ap}$	or $\dfrac{\text{their } \frac{dy}{dt}}{\text{their } \frac{dx}{dt}}$	
	When $x = ap^2$, $\dfrac{dy}{dx} = \dfrac{\sqrt{a}}{\sqrt{ap^2}} = \dfrac{\sqrt{a}}{\sqrt{a}\,p} = \dfrac{1}{p}$	$\dfrac{dy}{dx} = \dfrac{1}{p}$	A1
	or $\dfrac{dy}{dx} = \dfrac{4a}{2(2ap)} = \dfrac{1}{p}$		
	So $m_N = -p$	Applies $m_N = \dfrac{-1}{their\ m_T}$	M1
	N: $y - 2ap = -p\left(x - ap^2\right)$	Applies $y - 2ap = \left(\text{their } m_N\right)\left(x - ap^2\right)$	M1
	N: $y - 2ap = -px + ap^3$		
	N: $y + px = ap^3 + 2ap$	Correct solution.	A1 **cso** *
			[5]
(b)	$\left(6a, 0\right) \Rightarrow 0 + p\left(6a\right) = ap^3 + 2ap$	Substitutes $x = 6a$, $y = 0$ into **N**	M1
	$\Rightarrow 4ap = ap^3 \Rightarrow p = 2$	$p = 2$	A1
	$x = -a$, $p = 2 \Rightarrow y + 2\left(-a\right) = a(2)^3 + 2a(2)$	Substitutes $x = -a$ and their p into **N**	dM1
	$\Rightarrow y = 8a + 4a + 2a = 14a \Rightarrow D\left(-a, 14a\right)$	$D\left(-a, 14a\right)$	A1
			[4]
(c)	When $p = 2$, $x = a(2)^2 = 4a$	Substitutes their p into $x = ap^2$	M1
	$\text{Area}(XPD) = \dfrac{1}{2}\left(14a\right)\left(5a\right) = 35a^2$	Applies $\dfrac{1}{2}(\text{their } 14a)(\text{their "}4a\text{"} + a)$	M1
		$35a^2$	A1
			[3]
			12

Question Number	Scheme	Notes	Marks
9.	$(3-\mathrm{i})z^* + 2\mathrm{i}\,z = 9 - \mathrm{i}$		
	$(3-\mathrm{i})(x-\mathrm{i}y) + 2\mathrm{i}(x+\mathrm{i}y) = 9 - \mathrm{i}$	Substituting $z = x + \mathrm{i}y$ and $z^* = x - \mathrm{i}y$ into $(3-\mathrm{i})z^* + 2\mathrm{i}\,z = 9 - \mathrm{i}$	M1
	$\underline{3x - 3\mathrm{i}y - \mathrm{i}\,x - y} + 2\mathrm{i}\,x - 2y \ \{= 9 - \mathrm{i}\}$	Multiplies out $\underline{(3-\mathrm{i})(x-\mathrm{i}y)}$ correctly. This mark can be implied by correct later working.	A1
	Re part: $3x - y - 2y = 9$ Im part: $-3y - x + 2x = -1$ $3x - 3y = 9$ $x - 3y = -1$	Equating either real or imaginary parts. One set of correct equations. Correct equations.	M1 A1 A1
	$2x = 10 \Rightarrow x = 5$ $x - 3y = -1 \Rightarrow 5 - 3y = -1 \Rightarrow y = 2$	Attempt to solve simultaneous equations to find one of x or y. Either $x = 5$ or $y = 2$. Both $x = 5$ and $y = 2$.	ddM1 A1 A1
	$\{z = 5 + 2\mathrm{i}\}$		**[8]** **8**

Question Number	Scheme	Notes	Marks
10. (i)	$u_{n+1} = 5u_n + 3$, $u_1 = 3$ and $u_n = \frac{3}{4}(5^n - 1)$		
	$n = 1;\quad u_1 = \frac{3}{4}(5^1 - 1) = \frac{3}{4}(4) = 3$	Check that $u_n = \frac{3}{4}(5^n - 1)$ yields 3 when $n = 1$.	B1
	So u_n is true when $n = 1$.		
	Assume that for $n = k$ that, $u_k = \frac{3}{4}(5^k - 1)$ is true for $k \in \mathbb{Z}^+$.		
	Then $u_{k+1} = 5u_k + 3$		
	$= 5\left(\frac{3}{4}(5^k - 1)\right) + 3$	Substituting $u_k = \frac{3}{4}(5^k - 1)$ into $u_{k+1} = 5u_k + 3$	M1
	$= \frac{3}{4}(5)^{k+1} - \frac{15}{4} + 3$	An attempt to multiply out in order to achieve $\pm\lambda(5^{k+1}) \pm$ constant	M1
	$= \frac{3}{4}(5)^{k+1} - \frac{3}{4}$		
	$= \frac{3}{4}(5^{k+1} - 1)$	$\frac{3}{4}(5^{k+1} - 1)$	A1
	Therefore, the general statement, $u_n = \frac{3}{4}(5^n - 1)$ is true when $n = k + 1$. (As u_n is true for $n = 1$,) then u_n is true for all positive integers by mathematical induction	True when $n = k + 1$, then by induction the result is true for all positive integers.	A1
			[5]

Question Number	Scheme	Notes	Marks
10. (ii)	$f(n) = 5\left(5^n\right) - 4n - 5$ is divisible by 16		
	$f(1) = 5(5^1) - 4(1) - 5 = 16,$ {which is divisible by 16}.	Shows that $f(1) = 16$	B1
	{$\therefore f(n)$ is divisible by 16 when $n = 1$.}		
	Assume that for $n = k$,		
	$f(k) = 5\left(5^k\right) - 4k - 5$ is divisible by 16 for $k \in \mathbb{Z}^+$.		
	$f(k+1) - f(k) = 5\left(5^{k+1}\right) - 4(k+1) - 5 - \left(5\left(5^k\right) - 4k - 5\right)$	Applies $f(k+1) - f(k)$. Correct expression for $f(k+1) - f(k)$.	M1 A1
	$= 5\left(5^{k+1}\right) - 4k - 4 - 5 - 5\left(5^k\right) + 4k + 5$		
	$= 25\left(5^k\right) - 4k - 4 - 5 - 5\left(5^k\right) + 4k + 5$	Achieves an expression in 5^k.	M1
	$= 20\left(5^k\right) - 4$		
	$= 4\left(5\left(5^k\right) - 4k - 5\right) + 16k + 20 - 4$		
	$= 4\left(5\left(5^k\right) - 4k - 5\right) + 16k + 16$		
	$= 4f(k) + 16(k+1)$		
	$\therefore f(k+1) = 5f(k) + 16(k+1)$	$f(k+1) = 5f(k) + 16(k+1)$	A1
	{$\therefore f(k+1) = 5f(k) + 16(k+1)$, which is divisible by 16 as both $5f(k)$ and $16(k+1)$ are both divisible by 16.}		
	If the result is true for $n = k$, then it is now true for $n = k+1$. As the result has shown to be true for $n = 1$, then the result is true for all *n*.	Correct conclusion	A1 cso
			[6] **11**

Appendix

- dM1 denotes a method mark which is dependent upon the award of the previous method mark.
- ddM1 denotes a method mark which is dependent upon the award of the previous two method marks.
- ft denotes "follow through"
- cao denotes "correct answer only"
- oe denotes "or equivalent"

Other Possible Solutions

Question Number	Scheme	Notes	Marks
7.	$\mathbf{P} = \begin{pmatrix} 3a & -2a \\ -b & 2b \end{pmatrix}$, $\mathbf{M} = \begin{pmatrix} -6a & 7a \\ 2b & -b \end{pmatrix}$		
Aliter	$\mathbf{M} = \mathbf{PQ}$		
(b)	$\begin{pmatrix} -6a & 7a \\ 2b & -b \end{pmatrix} = \begin{pmatrix} 3a & -2a \\ -b & 2b \end{pmatrix}\begin{pmatrix} q_1 & q_2 \\ q_3 & q_4 \end{pmatrix}$		
Way 2	$-6 = 3q_1 - 2q_3$ $\quad 7 = 3q_2 - 2q_4$ $2 = -q_1 + 2q_3$, $\quad -1 = -q_2 + 2q_4$ $= \begin{pmatrix} -2 & 3 \\ 0 & 1 \end{pmatrix}$	Writes down a relevant pair of simultaneous equations. Can be implied by later working. Two out of four elements correct. Correct matrix.	M1 A1 A1 **[3]**

Question Number	Scheme	Notes	Marks
Aliter	$f(n) = 5\left(5^n\right) - 4n - 5$ is divisible by 16		
10. (ii)	$f(1) = 5(5^1) - 4(1) - 5 = 16,$	Shows that $f(1) = 16$	B1
Way 2	{which is divisible by 16}.		
	$\{\therefore f(n)$ is divisible by 16 when $n = 1.\}$		
	Assume that for $n = k$,		
	$f(k) = 5\left(5^k\right) - 4k - 5$ is divisible by 16 for $k \in \mathbb{Z}^+$.		
	$f(k+1) = 5\left(5^{k+1}\right) - 4(k+1) - 5$	Applies $f(k+1)$.	M1
		Correct expression for $f(k+1)$.	A1
	$= 25\left(5^k\right) - 4k - 9$	Achieves an expression in 5^k.	M1
	$= 5\left(5\left(5^k\right) - 4k - 5\right) + 20k + 25 - 4k - 9$		
	$= 5\left(5\left(5^k\right) - 4k - 5\right) + 16(k+1)$		
	$\therefore f(k+1) = 5f(k) + 16\left(k+1\right)$	$f(k+1) = 5f(k) + 16\left(k+1\right)$	A1
	$\{\therefore f(k+1) = 5f(k) + 16\left(k+1\right)$, which is divisible by 16		
	as both $5f(k)$ and $16\left(k+1\right)$ are both divisible by 16.$\}$		
	If the result is true for $n = k$, then it is now true for		
	$n = k+1$. As the result has shown to be true for $n = 1$,	Correct conclusion	A1 cso
	then the result is true for all *n*.		
			[6]

GradeMax

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE