C3 5164 UL

GradeMax

Mathematics ; 2014 ; May/Jun ; Paper 3 QP

1. The curve C has equation $y = f(x)$ where

$$f(x) = \frac{4x+1}{x-2}, \quad x > 2$$

(a) Show that

$$f'(x) = \frac{-9}{(x-2)^2}$$

(3)

Given that P is a point on C such that $f'(x) = -1$,

(b) find the coordinates of P.

(3)

a) $u = 4x+1$ $\quad v = x-2$ $\quad f'(x) = \dfrac{4(x-2)-(4x+1)}{(x-2)^2}$

$u' = 4$ $\quad v' = 1$

$= \dfrac{4x-8-4x-1}{(x-2)^2} = \dfrac{-9}{(x-2)^2}$ #

b) $f'(x) = -1 \Rightarrow (x-2)^2 = 9 \quad x = 2 \pm 3 \quad \therefore x = 5$

$f(5) = \dfrac{21}{3} = 7 \quad P(5,7)$

2. Find the exact solutions, in their simplest form, to the equations

Mathematics · 2014 · May/Jun · Paper 3 · QP

(a) $2\ln(2x + 1) - 10 = 0$

(2)

(b) $3^x e^{4x} = e^7$

(4)

a) $\ln(2x+1) = 5$

$2x+1 = e^5 \quad \therefore x = \frac{-1+e^5}{2}$

b) $3^x = \frac{e^7}{e^{4x}} \Rightarrow 3^x = e^{7-4x}$

$\ln 3^x = 7-4x$

$4x + x\ln 3 = 7$

$x(4+\ln 3) = 7 \quad \therefore x = \frac{7}{4+\ln 3}$

GradeMax

3. The curve C has equation $x = 8y \tan 2y$

The point P has coordinates $\left(\pi, \frac{\pi}{8}\right)$

(a) Verify that P lies on C.

(1)

(b) Find the equation of the tangent to C at P in the form $ay = x + b$, where the constants a and b are to be found in terms of π.

(7)

a) $y = \frac{\pi}{8}$ $\quad x = 8\left(\frac{\pi}{8}\right)\tan\left(\frac{\pi}{4}\right) = \pi \times 1 = \pi$ ✱

b) $u = 8y \quad v = \tan 2y$

$u' = 8 \quad v' = 2\sec^2 2y$

$$\frac{dx}{dy} = 8\tan 2y + 16y\sec^2 2y$$

$$\therefore \frac{dy}{dx} = \frac{1}{8\tan 2y + 16y\sec^2 2y}$$

$y = \frac{\pi}{8}$ $\quad M_t = \dfrac{1}{8\tan\left(\frac{\pi}{4}\right) + \dfrac{2\pi}{\left(\cos\frac{\pi}{4}\right)^2}} = \dfrac{1}{8 + 4\pi}$

$$y - \frac{\pi}{8} = \frac{1}{(8+4\pi)}(x - \pi) \Rightarrow (8+4\pi)y - \frac{\pi}{8}(8+4\pi) = x - \pi$$

$$\Rightarrow (8+4\pi)y = x - \pi + \pi + \frac{\pi^2}{2}$$

$$4(2+\pi)y = x - \frac{\pi^2}{2}$$




Figure 1

Figure 1 shows part of the graph with equation $y = f(x)$, $x \in \mathbb{R}$.

The graph consists of two line segments that meet at the point $Q(6, -1)$.

The graph crosses the y-axis at the point $P(0, 11)$.

Sketch, on separate diagrams, the graphs of

(a) $y = |f(x)|$ **(2)**


(b) $y = 2f(-x) + 3$ **(3)**

On each diagram, show the coordinates of the points corresponding to P and Q.


Given that $f(x) = a|x - b| - 1$, where a and b are constants,

(c) state the value of a and the value of b. **(2)**

a)



b)



c) $x=6, f(x)=-1 \quad a|6-b|-1=-1 \Rightarrow a|6-b|=0 \quad a=0, b=6$

$x=0 \quad f(x)=11 \quad a|-b|-1=11 \quad ax\pm b=12 \quad a\neq 0$

$b=6 \quad a=2$

GradeMax

5. $$g(x) = \frac{x}{x+3} + \frac{3(2x+1)}{x^2+x-6}, \quad x > 3$$

(a) Show that $g(x) = \frac{x+1}{x-2}, \quad x > 3$ **(4)**

(b) Find the range of g. **(2)**

(c) Find the exact value of a for which $g(a) = g^{-1}(a)$. **(4)**

a) $g(x) = \frac{x}{x+3} + \frac{3(2x+1)}{(x+3)(x-2)} = \frac{x(x-2)+3(2x+1)}{(x+3)(x-2)}$

$g(x) = \frac{x^2-2x+6x+3}{(x+3)(x-2)} = \frac{x^2+4x+3}{(x+3)(x-2)} = \frac{(x+3)(x+1)}{(x+3)(x-2)}$ ##

b) $g(x) > 1 \quad \frac{3+1}{3-2} = \frac{4}{1} = 4 \quad \therefore \underline{1 < g(x) < 4}$

c) $x = \frac{y+1}{y-2} \Rightarrow xy - 2x = y + 1 \Rightarrow xy - y = 1 + 2x$

$\therefore y = \frac{1+2x}{x-1} = g^{-1}(x)$

$\frac{a+1}{a-2} = \frac{1+2a}{a-1} \Rightarrow a^2 - 1 = 2a^2 - 3a - 2$


$\Rightarrow a^2 - 3a - 1 = 0$

$(a - \frac{3}{2})^2 - \frac{9}{4} = 1$

$(a - \frac{3}{2})^2 = \frac{13}{4}$

$a = \frac{3}{2} + \frac{\sqrt{13}}{2}$

$\frac{x+1}{x-2} \quad x \neq 2$






Figure 2

Figure 2 shows a sketch of part of the curve with equation

$$y = 2\cos\left(\frac{1}{2}x^2\right) + x^3 - 3x - 2$$

The curve crosses the x-axis at the point Q and has a minimum turning point at R.

(a) Show that the x coordinate of Q lies between 2.1 and 2.2

(2)

(b) Show that the x coordinate of R is a solution of the equation

$$x = \sqrt{1 + \frac{2}{3}x\sin\left(\frac{1}{2}x^2\right)}$$

(4)

Using the iterative formula

$$x_{n+1} = \sqrt{1 + \frac{2}{3}x_n\sin\left(\frac{1}{2}x_n^2\right)}, \quad x_0 = 1.3$$

(c) find the values of x_1 and x_2 to 3 decimal places.

(2)

a) $x = 2.1$ $y = -0.22$ ∴ by sign change rule
$x = 2.2$ $y = 0.55$ $2.1 < x < 2.2$

b) $x = \sqrt{1 + \frac{2}{3}x\sin\left(\frac{1}{2}x^2\right)}$ when $\frac{dy}{dx} = 0$ TP.

$\frac{dy}{dx} = -2x\sin\left(\frac{x^2}{2}\right) + 3x^2 - 3$ ∴ at TP $3x^2 = 3 + 2x\sin\left(\frac{x^2}{2}\right)$

$\therefore x^2 = 1 + \frac{2}{3}x\sin\left(\frac{x^2}{2}\right)$

c) $x_0 = 1.3$ $x_1 = 1.284$ $x_2 = 1.276$ $\therefore x = \sqrt{1 + \frac{2}{3}x\sin\left(\frac{x^2}{2}\right)}$

GradeMax

7. (a) Show that

$$\operatorname{cosec} 2x + \cot 2x = \cot x, \quad x \neq 90n°, \quad n \in \mathbb{Z}$$

(5)

(b) Hence, or otherwise, solve, for $0 \leqslant \theta < 180°$,

$$\operatorname{cosec}(4\theta + 10°) + \cot(4\theta + 10°) = \sqrt{3}$$

You must show your working.

(*Solutions based entirely on graphical or numerical methods are not acceptable.*)

(5)

a) $\frac{1}{\sin 2x} + \frac{\cos 2x}{\sin 2x} =$

$= \frac{1 + (2\cos^2 x - 1)}{2\sin x \cos x} = \frac{2\cos^2 x}{2\sin x \cos x} = \frac{\cos x}{\sin x} = \cot x$ ∎

b) $\operatorname{cosec} 2x + \cot 2x = \cot x$

$\operatorname{cosec}(4\theta + 10) + \cot(4\theta + 10) = \sqrt{3}$

$\therefore \cot x = \sqrt{3} \Rightarrow \tan x = \frac{1}{\sqrt{3}} \quad \therefore x = 30, 210, 390, 570$

$2x = 4\theta + 10 = 60, 420, 780, 1040$

$4\theta = \quad = 50, 410, 770, 1030$

$\therefore \theta = 12.5°, 102.5°$

Mathematics · 2014 · May/Jun · Paper 3 · QP GradeMax

8. A rare species of primrose is being studied. The population, P, of primroses at tir after the study started is modelled by the equation

$$P = \frac{800e^{0.1t}}{1 + 3e^{0.1t}}, \qquad t \geqslant 0, \quad t \in \mathbb{R}$$

(a) Calculate the number of primroses at the start of the study.

(2)

(b) Find the exact value of t when $P = 250$, giving your answer in the form $a \ln(b)$ where a and b are integers.

(4)

(c) Find the exact value of $\frac{dP}{dt}$ when $t = 10$. Give your answer in its simplest form.

(4)

(d) Explain why the population of primroses can never be 270

(1)

a) $t=0 \quad P = \frac{800}{1+3} = \frac{200}{2}$

b) $250 = \frac{800e^{0.1t}}{1+3e^{0.1t}} \Rightarrow 800e^{0.1t} = 250 + 750e^{0.1t}$

$\Rightarrow 50e^{0.1t} = 250 \Rightarrow e^{0.1t} = 5 \Rightarrow 0.1t = \ln 5$

$\therefore t = 10\ln 5$ (2)

c) $u = 800e^{0.1t} \quad v = 1+3e^{0.1t}$

$u' = 80e^{0.1t} \quad v' = 0.3e^{0.1t}$

$= \frac{(1+3e^{0.1t})80e^{0.1t} - (800e^{0.1t})(0.3e^{0.1t})}{(1+3e^{0.1t})^2}$

$t=10 \quad \frac{80e + 240e^2 - 240e^2}{(1+3e)^2} = \frac{80e}{(1+3e)^2}$

GradeMax

d) $P = \dfrac{800e^{0.1t}}{1+3e^{0.1t}} \div e^{0.1t} = \dfrac{800}{e^{-0.1t}+3}$

as $t \to \infty$ $P \to \dfrac{800}{3} \to 266.\dot{6}$

$\therefore$ Population can never reach 270

GradeMax
Mathematics · 2014 · May/Jun · Paper 3 · QP

9. (a) Express $2\sin\theta - 4\cos\theta$ in the form $R\sin(\theta - \alpha)$, where R and α are constants,

and $0 < \alpha < \frac{\pi}{2}$

Give the value of α to 3 decimal places.

(3)

$$H(\theta) = 4 + 5(2\sin 3\theta - 4\cos 3\theta)^2$$

Find

(b) (i) the maximum value of $H(\theta)$,

(ii) the smallest value of θ, for $0 \leqslant \theta < \pi$, at which this maximum value occurs.

(3)

Find

(c) (i) the minimum value of $H(\theta)$,

(ii) the largest value of θ, for $0 \leqslant \theta < \pi$, at which this minimum value occurs.

(3)

$R\sin(\theta-\alpha) = R\sin\theta\cos\alpha - R\cos\theta\sin\alpha$


$2\sin\theta \quad -4\cos\theta$

$\frac{R\sin\alpha}{R\cos\alpha} = \frac{4}{2} \Rightarrow \tan\alpha = 2 \quad \therefore \alpha = 1.107^c$

$R = \sqrt{4^2+2^2} \quad R = 2\sqrt{5}$

$\therefore 2\sqrt{5}\sin(\theta - 1.107)$

b) $\text{Max}(2\sin 3\theta - 4\cos 3\theta) = 2\sqrt{5}$



when $3\theta - 1.107 = \frac{\pi}{2} \quad \theta = 0.89264..$

$H_{max} = 4 + 5(2\sqrt{5})^2 = 4 + 100 = 104$ when $\theta = \frac{0.893}{2}$

c) $H_{min} = 4 + 5(0) = 4$ $\quad 0 \leq \pi < \pi$

when $3\theta - 1.107 = 0, \pi, 2\pi ...$

$3\theta = 1.107, \pi + 1.107, 2\pi + 1.107, 3\pi + 1.107$

$\therefore \text{max}\,\theta = 2.463$