GradeMax

FP3 UK June 2016 Model Answers

Kprime2

Leave blank

1.

$$\mathbf{A} = \begin{pmatrix} -2 & 1 & -3 \\ k & 1 & 3 \\ 2 & -1 & k \end{pmatrix}, \text{ where } k \text{ is a constant}$$

Given that the matrix A is singular, find the possible values of k.

(4)

1.

$$\det A = -2\begin{vmatrix} 1 & 3 \\ -1 & k \end{vmatrix} - \begin{vmatrix} k & 3 \\ 2 & k \end{vmatrix} - 3\begin{vmatrix} k & 1 \\ 2 & -1 \end{vmatrix}$$

$$= -2(k+3) - (k^2 - 6) - 3(-k-2)$$

$$= -2k - 6 - k^2 + 6 + 3k + 6$$

$$= -k^2 + k + 6 = 0$$

$$\therefore k^2 - k - 6 = 0$$

$$\therefore (k-3)(k+2) = 0$$

$$\therefore k = 3$$

$$k = -2$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

P 4 4 5 6 ... Scanned by CamScanner

GradeMax

Leave blank

2. The curve C has equation

$$y = \frac{x^2}{8} - \ln x, \quad 2 \leqslant x \leqslant 3$$

Find the length of the curve C giving your answer in the form $p + \ln q$, where p and q are rational numbers to be found.

(7)

2. $y = \frac{x^2}{8} - \ln x$

$\frac{dy}{dx} = \frac{x}{4} - \frac{1}{x} = \frac{x^2}{4x} - \frac{4}{4x}$

$= \frac{x^2-4}{4x}$

$\therefore \left(\frac{dy}{dx}\right)^2 = \frac{(x^2-4)^2}{16x^2}$

$\therefore 1 + \left(\frac{dy}{dx}\right)^2 = 1 + \frac{(x^2-4)^2}{16x^2} = \frac{16x^2 + x^4 - 8x^2 + 16}{16x^2}$

$= \frac{x^4 + 8x^2 + 16}{16x^2} = \frac{(x^2+4)^2}{16x^2}$

$\therefore \sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \frac{x^2+4}{4x}$ $\frac{x^2+4}{4x}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Scanned by CamScanner

GradeMax

$$\therefore S = \int_2^3 \sqrt{1+\left(\frac{dx}{dx}\right)^2}\, dx$$

$$= \int_2^3 \frac{x^2+4}{4x}\, dx$$

$$= \int_2^3 \frac{1}{4}x + \frac{1}{x}\, dx$$

$$= \left[\frac{1}{8}x^2 + \ln x\right]_2^3$$

$$= \frac{9}{8} + \ln 3 - \frac{1}{2} - \ln 2$$

$$= \frac{5}{8} + \ln\frac{3}{2}$$

Scanned by CamScanner

GradeMax

Leave blank

3. (a) Prove that

$$\frac{\mathrm{d}(\operatorname{arcoth} x)}{\mathrm{d}x} = \frac{1}{1-x^2}$$

(3)

Given that $y = (\operatorname{arcoth} x)^2$,

(b) show that

$$(1-x^2)\frac{\mathrm{d}^2y}{\mathrm{d}x^2} - 2x\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{k}{1-x^2}$$

where k is a constant to be determined.

(5)

3(a). Let $y = \operatorname{arcoth} x$

$\frac{\partial y}{\partial x} =$ $\therefore \coth y = x$

$\therefore \frac{\partial y}{\partial x}\left(-\operatorname{cosech}^2 y\right) = 1$

$\operatorname{cosech}^2 y \equiv \coth^2 y - 1$

$\therefore \frac{\partial y}{\partial x}\left(-(\coth^2 y - 1)\right) = 1$

$\coth y = x \Rightarrow \coth^2 y = x^2$

$\therefore \frac{\partial y}{\partial x}\left(-(x^2-1)\right) = 1$

$\therefore \frac{\partial y}{\partial x} = \frac{1}{-(x^2-1)} = \frac{1}{1-x^2}$

as required.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

P 4 4 5 6 6 A 0 6 3 2

Scanned by CamScanner

GradeMax

Leave blank

Question 3 continued

(b) $y = (\text{arcoth}\, x)^2$

$$\therefore \frac{\partial y}{\partial x} = \frac{2\,\text{arcoth}\, x}{1-x^2} \Rightarrow -2x\frac{\partial y}{\partial x} = \frac{-4x\,\text{arcoth}\, x}{1-x^2}$$

$$\therefore \frac{\partial^2 y}{\partial x^2} = \frac{(1-x^2)\left(\frac{2}{1-x^2}\right) - 2\,\text{arcoth}\, x\,(-2x)}{(1-x^2)^2}$$

$$= \frac{2 + 4x\,\text{arcoth}\, x}{(1-x^2)^2}$$

$$\Rightarrow (1-x^2)\frac{\partial^2 y}{\partial x^2} = (1-x^2)\left(\frac{2+4x\,\text{arcoth}\, x}{(1-x^2)^2}\right)$$

$$= \frac{2+4x\,\text{arcoth}\, x}{1-x^2}$$

$$\therefore \text{LHS} = (1-x^2)\frac{\partial^2 y}{\partial x^2} - 2x\frac{\partial y}{\partial x} = \frac{2+4x\,\text{arcoth}\, x}{1-x^2} - \frac{4x\,\text{arcoth}\, x}{1-x^2}$$

$$= \frac{2+4x\,\text{arcoth}\, x - 4x\,\text{arcoth}\, x}{1-x^2} = \frac{2}{1-x^2}$$

$K = 2$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Scanned by CamScanner

Further Mathematics · 2016 · May/Jun · FP3 · QP

GradeMax

Leave blank

4. (i) Find, without using a calculator.

$$\int_3^5 \frac{1}{\sqrt{15 + 2x - x^2}}\,dx$$

giving your answer as a multiple of π.

(5)

(ii)

(a) Show that

$$5\cosh x - 4\sinh x = \frac{e^{2x} + 9}{2e^x}$$

(3)

(b) Hence, using the substitution $u = e^x$ or otherwise, find

$$\int \frac{1}{5\cosh x - 4\sinh x}\,dx$$

(4)

4(i). $15+2x-x^2 = -(x^2-2x-15)$

$= -((x-1)^2-16)$

$= 16-(x-1)^2$

$\therefore \int_3^5 \frac{1}{\sqrt{16-(x-1)^2}}\,dx = \left[\arcsin\left(\frac{x-1}{4}\right)\right]_3^5$

$= \arcsin(1) - \arcsin\frac{1}{2}$

$= \frac{\pi}{2} - \frac{\pi}{6} = \frac{\pi}{3}$.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

P 4 4 5 6 6 A 0 1 0 3 2

Scanned by CamScanner

GradeMax

Leave blank

Question 4 continued

(ii) LHS $= 5\cosh x - 4\sinh x = \frac{5e^x + 5e^{-x}}{2} - \frac{4e^x - 4e^{-x}}{2}$

$= \frac{5e^x + 5e^{-x} - 4e^x + 4e^{-x}}{2}$

$= \frac{e^x + 9e^{-x}}{2} = \frac{(e^x + 9e^{-x})e^x}{2e^x}$

$= \frac{e^{2x} + 9}{2e^x} = \text{RHS}$ as required.

(b) $u = e^x \Rightarrow \frac{du}{dx} = e^x = u \Rightarrow dx = \frac{1}{u} \cdot du$

$\therefore \int \frac{1}{5\cosh x - 4\sinh x}\, dx = \int \frac{2e^x}{e^{2x} + 9}\, dx$

$= \int \frac{2u}{u^2 + 9} \cdot \frac{1}{u}\, du$

$= \int \frac{2}{u^2 + 9}\, du = \frac{2}{3}\arctan\left(\frac{u}{3}\right) + C$

$= \frac{2}{3}\arctan\left(\frac{e^x}{3}\right) + C$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Scanned by CamScanner

GradeMax

Leave blank

5. The hyperbola H has equation

$$\frac{x^2}{16} - \frac{y^2}{9} = 1$$

The point $P(4\sec\theta, 3\tan\theta)$, $0 < \theta < \frac{\pi}{2}$, lies on H.

(a) Show that an equation of the normal to H at the point P is

$$3y + 4x\sin\theta = 25\tan\theta$$

(5)

The line l is the directrix of H for which $x > 0$

The normal to H at P crosses the line l at the point Q. Given that $\theta = \frac{\pi}{4}$

(b) find the y coordinate of Q, giving your answer in the form $a + b\sqrt{2}$, where a and b are rational numbers to be found.

(6)

5(a). @ P, $\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{3\sec^2\theta}{4\sec\theta\tan\theta}$

$= \frac{3\sec\theta}{4\tan\theta} = \frac{3}{4\sin\theta}$

$\therefore$ gradient of normal $= -\frac{4\sin\theta}{3}$

$y - y_1 = m(x - x_1) \Rightarrow y - 4\sec\theta$

$y - 3\tan\theta = \frac{3}{4\sin\theta}(x - 4\sec\theta)$

$\therefore\ y - 3\tan\theta = -\frac{4\sin\theta}{3}(x - 4\sec\theta)$

$\therefore\ 3y - 9\tan\theta = -4\sin\theta\, x + 16\tan\theta$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Scanned by CamScanner

GradeMax

Question 5 continued


$\therefore 3y + 4x\sin\theta = 25\tan\theta$

as required.

(b) $a^2 = 16$, $b^2 = 9$ $\Big\}$ $9 = 16(e^2 - 1)$

$\therefore e = \frac{5}{4}$

$\theta = \frac{\pi}{4} \Rightarrow 3y + 2\sqrt{2}x = 25$



directrix $\frac{a}{e} = \frac{16}{5}$

$x_Q = \frac{16}{5}$

$\Rightarrow 3y + \frac{32}{5}\sqrt{2} = 25$

$\therefore y_Q = \frac{25}{3} - \frac{32}{15}\sqrt{2}$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Scanned by CamScanner

GradeMax

Leave blank

6. $$\mathbf{M} = \begin{pmatrix} p & -2 & 0 \\ -2 & 6 & -2 \\ 0 & -2 & q \end{pmatrix}$$

where p and q are constants.

Given that $\begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$ is an eigenvector of the matrix **M**,

(a) find the eigenvalue corresponding to this eigenvector. **(3)**

(b) find the value of p and the value of q. **(3)**

Given that 6 is another eigenvalue of **M**,

(c) find a corresponding eigenvector. **(2)**

Given that $\begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$ is a third eigenvector of **M** with eigenvalue 3

(d) find a matrix **P** and a diagonal matrix **D** such that

$$\mathbf{P}^{\mathrm{T}}\mathbf{M}\mathbf{P} = \mathbf{D}$$

(3)

6(a). $\begin{pmatrix} p & -2 & 0 \\ -2 & 6 & -2 \\ 0 & -2 & q \end{pmatrix}\begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} = \lambda\begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$

$\therefore \begin{pmatrix} 2p+4 \\ -18 \\ 4+q \end{pmatrix} = \begin{pmatrix} 2\lambda \\ -2\lambda \\ \lambda \end{pmatrix} \Rightarrow -18 = -2\lambda$

$\therefore \lambda = 9$

P 4 4 5 6 6 A 0 1 8 3 2

Scanned by CamScanner

Leave blank

Question 6 continued

(b) $2p+4=2\lambda \Rightarrow 2p+4=18$

$\therefore p=7$

$4+q=9 \Rightarrow q=5$

(c) $\begin{pmatrix} 7 & -2 & 0 \\ -2 & 6 & -2 \\ 0 & -2 & 5 \end{pmatrix}\begin{pmatrix} x \\ y \\ z \end{pmatrix} = 6\begin{pmatrix} x \\ y \\ z \end{pmatrix}$

$\therefore \begin{pmatrix} 7x-2y \\ -2x+6y-2z \\ -2y+5z \end{pmatrix} = \begin{pmatrix} 6x \\ 6y \\ 6z \end{pmatrix}$

$7x-2y=6x \Rightarrow x=2y$

$-2y+5z=6z \Rightarrow z=-2y$

$\Rightarrow x=2y=-z$

Let $x=2 \Rightarrow y=1 \Rightarrow z=-2$

$\therefore$ E-vector is $\begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Scanned by CamScanner

GradeMax

Question 6 continued

Leave blank

(d) $\lambda = 9$ & $\begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$

Normalise $\Rightarrow \frac{1}{3}\begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$

$\lambda = 6$ & $\begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix}$

Normalise $\Rightarrow \frac{1}{3}\begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix}$

$\lambda = 3$ & $\begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$ Normalise $\Rightarrow \frac{1}{3}\begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$

$\therefore P = \frac{1}{3}\begin{pmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & -2 & 2 \end{pmatrix}$

$D = \begin{pmatrix} 9 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 3 \end{pmatrix}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Scanned by CamScanner

GradeMax

Leave blank

7. Given that

$$I_n = \int \frac{\sin nx}{\sin x}\,dx, \quad n \geqslant 1$$

(a) prove that, for $n \geqslant 3$

$$I_n - I_{n-2} = \int 2\cos(n-1)x\,dx$$

(3)

(b) Hence, showing each step of your working, find the exact value of

$$\int_{\frac{\pi}{12}}^{\frac{\pi}{6}} \frac{\sin 5x}{\sin x}\,dx$$

giving your answer in the form $\frac{1}{12}(a\pi + b\sqrt{3} + c)$, where a, b and c are integers to be found.

(7)

$$I_n - I_{n-2} = \int \frac{\sin nx}{\sin x}\,dx - \int \frac{\sin[(n-2)x]}{\sin x}\,dx$$

$$= \int \frac{\sin(nx) - \sin(nx-2x)}{\sin x}\,dx$$

$$= \int \frac{2\cos\left(\frac{2nx-2x}{2}\right)\sin\left(\frac{2x}{2}\right)}{\sin x}\,dx \quad *$$

$$= \int 2\cos(n-1)x\,dx$$

* Using $\sin A - \sin B \equiv 2\cos\left(\frac{A+B}{2}\right)\sin\left(\frac{A-B}{2}\right)$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Scanned by CamScanner

GradeMax

Leave blank

Question 7 continued

$$I_5 - I_3 = 2\int_{\pi/12}^{\pi/6} \cos 4x \, dx$$

$$= 2\left[\tfrac{1}{4}\sin 4x\right]_{\pi/12}^{\pi/6}$$

$$= 0 \Rightarrow I_5 = I_3$$

$$I_3 - I_1 = \int_{\pi/12}^{\pi/6} 2\cos 2x \, dx$$

$$= \left[\sin 2x\right]_{\pi/12}^{\pi/6}$$

$$= \frac{-1+\sqrt{3}}{2}$$

$$I_1 = \int_{\pi/12}^{\pi/6} 1 \, dx = \frac{\pi}{6} - \frac{\pi}{12} = \frac{\pi}{12}$$

$$\therefore I_3 = \frac{\pi}{12} + \frac{-1+\sqrt{3}}{2} = \frac{\pi}{12} + \frac{-6+6\sqrt{3}}{12}$$

$$\Rightarrow I_5 = \frac{1}{12}(\pi + 6\sqrt{3} - 6) \qquad a=1,\ b=6,\ c=-6$$

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

P 4 4 5 6 6 A 0 2 6 3 2
Scanned by CamScanner

GradeMax

Leave blank

8. The plane Π_1 has equation

$$x - 5y - 2z = 3$$

The plane Π_2 has equation

$$\mathbf{r} = \mathbf{i} + 2\mathbf{j} + \mathbf{k} + \lambda(\mathbf{i} + 4\mathbf{j} + 3\mathbf{k}) + \mu(2\mathbf{i} - \mathbf{j} + \mathbf{k})$$

where λ and μ are scalar parameters.

(a) Show that Π_1 is perpendicular to Π_2

(4)

(b) Find a cartesian equation for Π_2

(2)

(c) Find an equation for the line of intersection of Π_1 and Π_2 giving your answer in the form $(\mathbf{r} - \mathbf{a}) \times \mathbf{b} = \mathbf{0}$, where $\mathbf{a}$ and $\mathbf{b}$ are constant vectors to be found.

(6)

8(a) Π_2: $\underset{\sim}{n}_2 = \begin{pmatrix}1\\4\\3\end{pmatrix} \times \begin{pmatrix}2\\-1\\1\end{pmatrix} = \begin{matrix}1 & 4 & 3 & 1 & 4\\ 2 & -1 & 1 & 2 & -1\end{matrix} \times \begin{matrix}3\\1\end{matrix}$

$= \begin{pmatrix}7\\5\\-9\end{pmatrix}$

If π_1 & π_2 are perpendicular then their normals must be perpendicular.

Π_1: $\underset{\sim}{n}_1 = \begin{pmatrix}1\\-5\\-2\end{pmatrix}$

$\underset{\sim}{n}_1 \cdot \underset{\sim}{n}_2 = \begin{pmatrix}7\\5\\-9\end{pmatrix} \cdot \begin{pmatrix}1\\-5\\-2\end{pmatrix} = 7 - 25 + 18 = 0$

$\Rightarrow$ π_1 & π_2 are perpendicular.

Scanned by CamScanner

DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA

GradeMax

Leave blank

Question 8 continued

(b) $L \cdot \begin{pmatrix} 7 \\ 5 \\ -9 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 7 \\ 5 \\ -9 \end{pmatrix}$

$= 8$

$\Rightarrow 7x + 5y - 9z = 8$

(c) $x - 5y - 2z = 3 \Rightarrow 5y = x - 2z - 3$

$7x + 5y - 9z = 8$

$\therefore 7x + x - 2z - 3 - 9z = 8$

$\therefore 8x - 11z = 11$

$\therefore x = \frac{11}{8} + \frac{11}{8}z$

$\therefore 5y = \frac{11}{8} + \frac{11}{8}z - 2z - 3$

$5y = -\frac{13}{8} - \frac{5}{8}z$

$\therefore y = -\frac{13}{40} - \frac{1}{8}z$

$z = z$

$\therefore \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \frac{11}{8} \\ -\frac{13}{40} \\ 0 \end{pmatrix} + z \begin{pmatrix} 11/8 \\ -1/8 \\ 1 \end{pmatrix}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Scanned by CamScanner

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Leave blank

Question 8 continued

$$= \begin{pmatrix} 11/8 \\ -13/40 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 11 \\ -1 \\ 8 \end{pmatrix}$$

$$\therefore \left(\underset{\sim}{r} - \begin{pmatrix} \frac{11}{8} \\ -13/40 \\ 0 \end{pmatrix} \right) \times \begin{pmatrix} 11 \\ -1 \\ 8 \end{pmatrix} = 0$$

Note: There are many possible solutions for this part of the question.

Solutions will vary depending on what you expressed your x, y, z in terms of...

Scanned by CamScanner